arXiv:1603.01283v2 [math.NT] 16 Mar 2016

ABELIAN PART OF COMPATIBLE SYSTEM AND GEOMETRY OF ROOTS

Chun Yin Hui

Mathematics Research Unit, University of Luxembourg,
6 rue Richard Coudenhove-Kalergi, L-1359 Luxembourg
Email: pslnfq@gmail.com

Abstract. Let $\{V_\ell\}_\ell$ be a strictly compatible rational system of Galois representations arising from geometry. Let V_ℓ^{ss} be the semisimplification of V_ℓ, V_ℓ^{ab} the maximal abelian subrepresentation of V_ℓ^{ss}, and $\mathbf{G}_\ell$ the algebraic monodromy group of V_ℓ^{ss}. Conjecturally, the abelian part $\{V_\ell^{\mathrm{ab}}\}_\ell$ is again a strictly compatible rational system (Conj. Ab). This conjecture has meaningful consequences to arithmetic geometry, for example, the Tate conjecture is independent of ℓ if Conj. Ab and the semisimplicity conjecture of Grothendieck hold. The main theme of this paper is to investigate Conj. Ab via ℓ-independence of $\mathbf{G}_\ell$. More precisely, we prove that Conj. Ab follows if (i) $\mathbf{G}_\ell$ is connected for all ℓ, (ii) the tautological representation of the complex semisimple Lie group $\mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C})$ is independent of ℓ, and (iii) the roots of $\mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C})$ meet a criterion (Thm. 1.1). These conditions are satisfied under some group theoretic assumptions on $\mathbf{G}_\ell$ (Thm. 1.3), for example, (ii) and (iii) hold if $\mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C})$ is connected and semisimple of type $A_4 \times A_6 \times A_6$ for some ℓ; (iii) holds if $\mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C})$ is almost simple of type different from A_7, A_8, B_4, D_8.

Date: 16th March, 2016.

2010 *Mathematics Subject Classification.* 11F80, 14F20, 20G30 (primary), 14C15, 17B20 (secondary).

Key words and phrases. Galois representations, motives, ℓ-independence, algebraic monodromy groups, strictly compatible system, Frobenius tori, locally algebraic abelian representations, formal bi-character, complex semisimple Lie groups, root systems.

The present project is supported by the National Research Fund, Luxembourg, and cofunded under the Marie Curie Actions of the European Commission (FP7-COFUND)..

1. Introduction

1.1. **Motivation.** Fix a number field K and an embedding $\bar{\mathbb{Q}}_{\ell'} \hookrightarrow \mathbb{C}$. The Langlands reciprocity conjecture [Lan79, §2] predicts a one-to-one correspondence between the isomorphism classes of irreducible cuspidal automorphic representations π of $\mathrm{GL}_n(\mathbb{A}_K)$ and the isomorphism classes of n-dimensional irreducible $\bar{\mathbb{Q}}_{\ell'}$-representations $\bar{W}_{\ell'}$ of $\mathrm{Gal}(\bar{K}/K)$ *arising from geometry*, which preserves the local L-factors at almost all primes p: if π corresponds to $\bar{W}_{\ell'}$, then $L_p(\pi, p^{-s}) = L_p(\bar{W}_{\ell'} \otimes \mathbb{C}, p^{-s})$. Arising from geometry here means that $\bar{W}_{\ell'}$ is isomorphic to a subquotient of $\bigoplus_{j=1}^{m} H^{i_j}(\bar{X}_j, \bar{\mathbb{Q}}_{\ell'}(r_j))$, where X_j is smooth projective over K, $\bar{X}_j := X_j \times \bar{K}$, $m \in \mathbb{N}$, $1 \leq j \leq m$, and $i_j, r_j \in \mathbb{Z}$. For the complete description of the one-to-one correspondence, see [Cl90, §4],[Tay04, Conj. 1.1, 3.4, 3.5].

Let $\pi \leftrightarrow \bar{W}_{\ell'}$ be a correspondence arising from the Langlands reciprocity conjecture as above. Since the automorphic side of the conjecture is independent of ℓ, one obtains a correspondence $\pi \leftrightarrow \bar{W}_\ell$ for every prime ℓ by the conjecture at ℓ and thus a system of $\bar{\mathbb{Q}}_\ell$-representations $\{\bar{W}_\ell\}_\ell$ of $\mathrm{Gal}(\bar{K}/K)$. Since every $\bar{W}_\ell$ arises from geometry and the local L-factors are preserved, the system $\{\bar{W}_\ell\}_\ell$ satisfies *weak compatibility* in the following sense:

Definition 1. Let $\rho_\iota : \mathrm{Gal}(\bar{K}/K) \to \mathrm{GL}_n(F_\iota)$ be a Galois representation over a characteristic zero field F_ι for every ι belonging to an index set I. The system $\{\rho_\iota\}_\iota$ is said to be *weakly compatible* if there exists an embedding $F_\iota \hookrightarrow \mathbb{C}$ for every ι such that:

(i) for each ι, ρ_ι is unramified at almost all finite places v of K;
(ii) for every pair ι, ι' of indices, the characteristic polynomials $P_{v,\rho_\iota}(t), P_{v,\rho_{\iota'}}(t)$ of the Frobenii above v agree in $\mathbb{C}[t]$ via $F_\iota \hookrightarrow \mathbb{C} \hookleftarrow F_{\iota'}$ for almost all finite places v.

Let X be smooth projective over K. The Galois representation $H^i(\bar{X}, \mathbb{Q}_\ell)$ is conjectured to be semisimple (Grothendieck). Deligne has proven that the system of ℓ-adic representations $\{V_\ell = H^i(\bar{X}, \mathbb{Q}_\ell)\}_\ell$ is *strictly compatible rational* (see §2.2) [De74]. The semisimplicity conjecture, the conjectural correspondence of Langlands, and the compatibility condition lead us to the following conjecture on the decomposition of the *strictly compatible rational system* $\{V_\ell\}_\ell$ *arising from geometry* (meaning that $\{V_\ell\}_\ell$ is a strictly compatible rational system and $V_\ell \otimes \bar{\mathbb{Q}}_\ell$ arises from geometry for all ℓ).

Conjecture De. Let $\{V_\ell\}_\ell$ be a strictly compatible rational system arising from geometry. Then there exist weakly compatible systems of $\bar{\mathbb{Q}}_\ell$-representations $\{\bar{W}_\ell^{(k)}\}_\ell$ for $1 \leq k \leq m$ such that for all ℓ, V_ℓ admits the following decomposition:

$$V_\ell \otimes \bar{\mathbb{Q}}_\ell \cong \bar{W}_\ell^{(1)} \oplus \bar{W}_\ell^{(2)} \oplus \cdots \oplus \bar{W}_\ell^{(m)}.$$

Since the Langlands reciprocity conjecture is wide open[1], Conjecture De in general is out of reach, especially when the irreducibles of $V_\ell \otimes \bar{\mathbb{Q}}_\ell$ are not all of dimension one or two. However, the next conjecture, which follows from the above one, is the motivation of this paper.

Conjecture Ab. Let $\{V_\ell\}_\ell$ be a strictly compatible rational system arising from geometry, with the exceptional set S. Denote by V_ℓ^{ab} the maximal abelian subrepresentation of the

[1] The function field analogue of the Langlands conjecture (for GL_n) is proved by Lafforgue [Laf02]. For recent advance of the Langlands conjecture, see [BLGGT14].

semisimplification V_ℓ^{ss} of V_ℓ. Then the abelian part $\{V_\ell^{\mathrm{ab}}\}_\ell$ is also strictly compatible rational with the same exceptional set S.

By Conjecture Ab, the abelian part $\{V_\ell^{\mathrm{ab}}\}_\ell$ is associated to some $\mathbb{Q}$-representation of some Serre group $\mathbf{S}_{\mathfrak{m}}$ (see §2.2). Conjecture Ab has the following meaningful consequences.

(i) *Motives and new compatible systems.* Let V_ℓ^{c} be the unique complementary representation of V_ℓ^{ab} in V_ℓ^{ss}. Then $\{V_\ell^{\mathrm{c}}\}_\ell$ is also strictly compatible rational. Hence, one should decompose $\{V_\ell^{\mathrm{ss}}\}_\ell$ into $\{V_\ell^{\mathrm{ab}}\}_\ell$ and $\{V_\ell^{\mathrm{c}}\}_\ell$ and study them separately. Actually, it is believed that $\{V_\ell^{\mathrm{ss}}\}_\ell$ is the realization of a *pure motive* M [Kl72] and the equation

$$\{V_\ell^{\mathrm{ab}}\}_\ell \oplus \{V_\ell^{\mathrm{c}}\}_\ell = \{V_\ell^{\mathrm{ss}}\}_\ell$$

corresponds to a decomposition of M. Let $\{V_\ell\}_\ell$, $\{W_\ell\}_\ell$, $\{U_\ell\}_\ell$, ... be strictly compatible rational systems arising from geometry. Then any tensor construction of them is still strictly compatible rational arising from geometry. Therefore, one produces new strictly compatible rational systems by Conjecture Ab, e.g., $\{(V_\ell \otimes W_\ell)^?\}_\ell$, $\{(\mathrm{Sym}^3 V_\ell)^?\}_\ell$, where $? \in \{\mathrm{ab}, \mathrm{c}\}$.

(ii) *Factorization of L-function.* Let $\rho := \{\rho_\ell : \mathrm{Gal}(\bar{K}/K) \to \mathrm{GL}(V_\ell)\}_\ell$ be a strictly compatible rational system arising from geometry and S the exceptional subset of finite places of K for ρ. Denote by $P_{v,\rho}(t) := \det(1 - \rho_\ell(\mathrm{Frob}_{\bar{v}})t) \in \mathbb{Q}[t]$ the characteristic polynomial (independent of ℓ, $\bar{v}$) of the Frobenius element at $\bar{v}$ on V_ℓ whenever $\bar{v}$ is a place of $\bar{K}$ extending v of K, $v \notin S$, and $v \nmid \ell$. The L-function of ρ (depends on S) is defined [Se98, Ch. I §2.5] to be

$$L(\rho, s) := \prod_{v \notin S} \frac{1}{P_{v,\rho}((Nv)^{-s})},$$

where $Nv \in \mathbb{N}$ is the norm of v. The L-function $L(\rho, s)$ converges in some right half plane $\mathrm{Re}(s) > C$. Denote by ρ^{ss} the semisimplification of the system ρ and by ρ^{ab} and ρ^{c} the abelian and complementary subsystems of ρ^{ss}. Since $L(\rho, s) = L(\rho^{\mathrm{ss}}, s)$ and ρ^{ab} and ρ^{c} are strictly compatible rational with the exceptional set S by Conjecture Ab, we obtain a factorization

$$L(\rho, s) = L(\rho^{\mathrm{ab}}, s) L(\rho^{\mathrm{c}}, s).$$

Since ρ^{ab} arises from CM motives (Deligne's absolute Hodge cycles) [DMOS82],[FM93, §6 Prop.], [Sc94], $L(\rho^{\mathrm{ab}}, s)$ has meromorphic continuation to the complex plane.

(iii) *ℓ-independence of the Tate conjecture.* Let X be a smooth projective variety defined over K, $\mathfrak{g}_\ell$ the Lie algebra of the image of $\mathrm{Gal}(\bar{K}/K)$ in $\mathrm{GL}(H^{2r}(\bar{X}, \mathbb{Q}_\ell(r)))$, and $CH^r(\bar{X})$ the Chow group of codimension r cycles of $\bar{X}$. The Tate conjecture [Ta65] asserts that the image of the cycle map

$$c_\ell : CH^r(\bar{X}) \to H^{2r}(\bar{X}, \mathbb{Q}_\ell(r))$$

generates the Lie algebra invariants $H^{2r}(\bar{X},\mathbb{Q}_\ell(r))^{\mathfrak{g}_\ell}$. Embed $\bar{K}$ in $\mathbb{C}$. The diagram

$$\begin{array}{ccc} CH^r(\bar{X}) & \xrightarrow{c_\ell} & H^{2r}(\bar{X},\mathbb{Q}_\ell(r)) \\ \downarrow & & \downarrow \alpha_\ell \\ CH^r(X(\mathbb{C})) & \xrightarrow{c_{\mathbb{C}}\otimes 1} & H^{2r}(X(\mathbb{C}),\mathbb{Q})\otimes\mathbb{Q}_\ell \end{array}$$

is commutative, where α_ℓ is the comparison isomorphism between étale and singular cohomology and $c_{\mathbb{C}}$ is the cycle map for the complex manifold $X(\mathbb{C})$. Assume $V_\ell = H^{2r}(\bar{X},\mathbb{Q}_\ell(r))$ is semisimple for all ℓ (e.g., X abelian variety [Fa83]), then $V_\ell^{\mathfrak{g}_\ell} = (V_\ell^{\mathrm{ab}})^{\mathfrak{g}_\ell}$. There exists a finite extension K' of K such that $(V_\ell^{\mathrm{ab}})^{\mathfrak{g}_\ell} = (V_\ell^{\mathrm{ab}})^{\mathrm{Gal}(\bar{K}/K')}$ for all ℓ. Since $\mathrm{Im}c_\ell \subset V_\ell^{\mathfrak{g}_\ell}$ and the dimension of $(V_\ell^{\mathrm{ab}})^{\mathrm{Gal}(\bar{K}/K')}$ (resp. $\mathbb{Q}_\ell\cdot\mathrm{Im}c_\ell$) is independent of ℓ by Conjecture Ab (resp. the diagram), we conclude that the Tate conjecture is independent of ℓ. Note that we only need Conjecture Ab for a big enough extension K' of K.

1.2. **Results.** Let $\{V_\ell\}_\ell$ be a strictly compatible rational system (§2.2) of ℓ-adic ($\mathbb{Q}_\ell$-) representations of $\mathrm{Gal}(\bar{K}/K)$ arising from geometry (§1.1). The semisimplification of V_ℓ is represented by

$$\varphi_\ell : \mathrm{Gal}(\bar{K}/K) \to \mathrm{GL}(V_\ell^{\mathrm{ss}}) \cong \mathrm{GL}_N(\mathbb{Q}_\ell)$$

for all ℓ. The *algebraic monodromy group* of V_ℓ^{ss}, denoted by $\mathbf{G}_\ell$, is defined as the Zariski closure of the Galois image $\varphi_\ell(\mathrm{Gal}(\bar{K}/K))$ in the $\mathbb{Q}_\ell$-algebraic group $\mathrm{GL}_{V_\ell^{\mathrm{ss}}}$. Since φ_ℓ is semisimple, we identify $\mathbf{G}_\ell$ as a reductive subgroup of $\mathrm{GL}_{N,\mathbb{Q}_\ell}$ for all ℓ. The tautological representation $\mathbf{G}_\ell^\circ \hookrightarrow \mathrm{GL}_{N,\mathbb{Q}_\ell}$ is conjectured to be independent of ℓ [Se94]. For results on ℓ-independence of monodromy group of compatible system, see [Se72, Se85],[Ri76],[Chi92],[Pi98],[LP95] for abelian varieties and [LP92],[Chin04],[Hui13, Hui15a, Hui15b] for more general case.

Suppose $\mathbf{G}_\ell$ is connected for all ℓ. The derived group $\mathbf{G}_\ell^{\mathrm{der}}$ of $\mathbf{G}_\ell$ is connected and semisimple. Embed $\mathbb{Q}_\ell$ into $\mathbb{C}$ for all ℓ. Let $\mathbf{T}_{\mathbb{C}}^{\mathrm{ss}}$ be a maximal torus of the complex semisimple Lie group $\mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C})$ and Γ the character group of $\mathbf{T}_{\mathbb{C}}^{\mathrm{ss}}$. Then the automorphism group of $\mathbf{T}_{\mathbb{C}}^{\mathrm{ss}}$ is canonically isomorphic to the automorphism group of Γ. The *roots* R of $\mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C})$ (with respect to $\mathbf{T}_{\mathbb{C}}^{\mathrm{ss}}$) are the subset of Γ given by the non-zero characters of the adjoint representation. The main theme of the paper is

$$\begin{array}{ccc} \ell\text{-independence of } \mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C}) \hookrightarrow \mathrm{GL}_N(\mathbb{C}) & & \\ + & \Rightarrow & \text{Conjecture Ab} \\ \text{a criterion on } R & & \end{array}$$

Theorem 1.1. *Let $\{V_\ell\}_\ell$ be a strictly compatible rational system arising from geometry, with the exceptional set S. Let $\mathbf{G}_\ell$ be the algebraic monodromy group of V_ℓ^{ss} for all ℓ. Suppose the following conditions hold:*

(i) $\mathbf{G}_\ell$ is connected for all ℓ;
(ii) the conjugacy class of $\mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C})$ in $\mathrm{GL}_N(\mathbb{C})$ is independent of ℓ;
(iii) the roots R of $\mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C})$ are stable under the normalizer $N_{\mathrm{GL}_N(\mathbb{C})}(\mathbf{T}_{\mathbb{C}}^{\mathrm{ss}})$.

Then the abelian part $\{V_\ell^{\mathrm{ab}}\}_\ell$ is strictly compatible rational with the same exceptional set S.

Remark 1.2. *Condition 1.1(i) is equivalent to $\mathbf{G}_{\ell'}$ being connected for some ℓ' by Serre (see §2.1). Thus, one can take a finite extension of K for 1.1(i) to hold. Condition 1.1(ii) implies that 1.1(iii) is independent of ℓ.*

The ℓ-independence condition 1.1(ii) and the roots criterion 1.1 (iii) can be checked in specific cases. However, they are always satisfied under *Hypothesis A* [Hui15b, §3] (see 1.3(ii)). Together with a determination of almost simple Lie groups satisfying these conditions, we obtain the next result.

Theorem 1.3. *Let $\{V_\ell\}_\ell$ be a strictly compatible rational system arising from geometry, with the exceptional set S. Let $\mathbf{G}_\ell$ be the algebraic monodromy group of V_ℓ^{ss} for all ℓ. Suppose (i) $\mathbf{G}_\ell$ is connected for all ℓ and one of the following conditions hold:*

(ii) Hypothesis A: there exists some ℓ' such that $\mathbf{G}_{\ell'}^{\mathrm{der}}(\mathbb{C})$ has at most one factor of type A_4 and if $\mathbf{H}_{\mathbb{C}}$ is an almost simple factor of $\mathbf{G}_{\ell'}^{\mathrm{der}}(\mathbb{C})$, then $\mathbf{H}_{\mathbb{C}}$ is of type A_n for some $n \in \mathbb{N}\backslash\{1,2,3,5,7,8\}$;

(ii') the conjugacy class of $\mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C})$ in $\mathrm{GL}_N(\mathbb{C})$ is independent of ℓ and $\mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C})$ is almost simple of type different from A_7, A_8, B_4, D_8.

Then the abelian part $\{V_\ell^{\mathrm{ab}}\}_\ell$ is strictly compatible rational with the same exceptional set S.

Remark 1.4. *The consequences of Conjecture Ab in* §1.1 *are valid under the group theoretic assumptions in Theorem 1.1 or 1.3.*

Under Hypothesis A, one can prove the following for Conjecture De.

Theorem 1.5. *Let $\{V_\ell\}_\ell$ be a strictly compatible rational system arising from geometry. Let $\mathbf{G}_\ell$ be the algebraic monodromy group of V_ℓ^{ss} for all ℓ. Suppose (i) $\mathbf{G}_\ell$ is connected for all ℓ and (ii) Hypothesis A is satisfied. Let $\bigoplus^m \bar{W}_{\ell'}$ be a $\bar{\mathbb{Q}}_{\ell'}$-subrepresentation of $V_{\ell'}^{\mathrm{ss}} \otimes \bar{\mathbb{Q}}_{\ell'}$ for some ℓ' and $\{\bar{W}_\ell\}_\ell$ a weakly compatible system of irreducible $\bar{\mathbb{Q}}_\ell$-representations satisfying the following conditions:*

(iii) $\{\bar{W}_\ell\}_\ell$ extends $\bar{W}_{\ell'}$;

(iv) for all ℓ, descend $\bar{W}_\ell$ to an E_ℓ-representation W_ℓ in which E_ℓ is a finite extension of $\mathbb{Q}_\ell$. Then the local representations of W_ℓ above ℓ are of Hodge-Tate type.

Then $\{\bigoplus^m \bar{W}_\ell\}_\ell$ is a subsystem of $\{V_\ell^{\mathrm{ss}} \otimes \bar{\mathbb{Q}}_\ell\}_\ell$.

1.3. **Outline.** The idea behind Theorems 1.1 and 1.5 is that an irreducible representation $\bar{W}_{\ell'}$ is a subrepresentation of a semisimple $\bar{V}_{\ell'}$ (denoted by $\bar{W}_{\ell'} \leq \bar{V}_{\ell'}$) if and only if $\bar{V}_{\ell'} \otimes \bar{W}_{\ell'}^* = \mathrm{Hom}(\bar{W}_{\ell'}, \bar{V}_{\ell'})$ has non-trivial invariants; to do the same thing for a family $\{\bar{W}_\ell, \bar{V}_\ell\}_\ell$ assuming $\bar{W}_{\ell'} \leq \bar{V}_{\ell'}$ for some ℓ', it suffices to show that the tautological representation $\bar{V}_\ell \otimes \bar{W}_\ell^*$ of its algebraic monodromy group $\mathbf{H}_\ell$ is independent of ℓ. Section 2 is devoted to the proofs of Theorems 1.1, (part of) 1.3, 1.5, which rely on the ideas we have developed in [Hui13, Hui15b]. The criterion on the roots R (1.1(iii)) ensures the desired ℓ-independence of the tautological representations of $\mathbf{H}_\ell$. Section 3 is devoted to the study of the criterion on R by a geometric configuration of $R \subset \Gamma \otimes \mathbb{R}$. The results will complete the proof of Theorem 1.3.

2. Abelian part of compatible system

2.1. **Frobenius tori.** Let E_ℓ be a finite extension of $\mathbb{Q}_\ell$ and fix an embedding $E_\ell \hookrightarrow \mathbb{C}$ for all ℓ. Let $\{V_\ell\}_\ell$ be a weakly compatible (Definition 1) system of semisimple E_ℓ-representations:

$$\psi_\ell : \mathrm{Gal}(\bar{K}/K) \to \mathrm{GL}(V_\ell), \quad \forall \ell. \tag{1}$$

The algebraic monodromy group $\mathbf{G}_\ell$ of V_ℓ is reductive. The method of *Frobenius tori*, pioneered by Serre, is foundational to the study of ℓ-independence of $\mathbf{G}_\ell$. Denote by v a finite

place of K and $\bar{v}$ a place of $\bar{K}$ dividing v. Let $\psi_\ell(\mathrm{Frob}_{\bar{v}})$ be the image of the Frobenius at $\bar{v}$ whenever ψ_ℓ is unramified at v.

Definition 2. The *Frobenius torus* $\mathbf{T}_{\bar{v},\ell}$ is defined as the identity component of the smallest algebraic subgroup of $\mathbf{G}_\ell$ containing the semisimple part of $\psi_\ell(\mathrm{Frob}_{\bar{v}})$.

Theorem 2.1. *(Serre* [Se81]*) (see also* [Chi92, Theorem 3.7],[LP97, Theorem 1.2]*) Fix a prime ℓ' and suppose $\mathbf{G}_{\ell'}$ is connected. Denote by p_v the characteristic of v and by q_v the cardinality of the residue field of v. Suppose the eigenvalues α of $\psi_{\ell'}(\mathrm{Frob}_{\bar{v}})$ are algebraic for almost all v and the following conditions hold:*

(i) the absolute values of α in all complex embeddings are equal;
(ii) α is a unit at any finite place not above p_v;
(iii) for any non-Archimedean valuation w of $\bar{\mathbb{Q}}$ such that $w(p_v) > 0$, the ratio $w(\alpha)/w(q_v)$ belongs to a finite subset of $\mathbb{Q}$ that is independent of $\bar{v}$.

Then there exists a proper closed subvariety $\mathbf{Y}$ of $\mathbf{G}_{\ell'}$ such that $\mathbf{T}_{\bar{v},\ell'}$ is a maximal torus of $\mathbf{G}_{\ell'}$ whenever $\psi_{\ell'}(\mathrm{Frob}_{\bar{v}}) \in \mathbf{G}_{\ell'} \backslash \mathbf{Y}$.

The following consequences are well-known.

Corollary 2.2. *Suppose the hypotheses of Theorem 2.1 hold. Then $\mathbf{G}_\ell$ is connected for all ℓ.*

Proof. By the Chebotarev density theorem, $\mathbf{G}_\ell$ is the Zariski closure of the image of the Frobenii $\psi_\ell(\mathrm{Frob}_{\bar{v}})$ for a density 1 subset S of places v. Moreover, we can choose S such that $\mathbf{T}_{\bar{v},\ell'}$ is a maximal torus of $\mathbf{G}_{\ell'}$ for all $v \in S$ by Theorem 2.1. Since $\mathbf{G}_{\ell'}$ is connected reductive, the smallest algebraic subgroup of $\mathbf{G}_{\ell'}$ containing $\psi_{\ell'}(\mathrm{Frob}_{\bar{v}})$ is connected if $\mathbf{T}_{\bar{v},\ell'}$ is a maximal torus. Hence, $\mathbf{G}_\ell$ is generated by the Frobenius tori $\mathbf{T}_{\bar{v},\ell}$ (for all $v \in S$) by the weak compatibility. We conclude that $\mathbf{G}_\ell$ is connected for all ℓ. □

Corollary 2.3. *Suppose the hypotheses of Theorem 2.1 hold. Let $\mathbf{T}_{\ell,\mathbb{C}}$ be a maximal torus of $\mathbf{G}_\ell(\mathbb{C})$. Then the tautological representation of $\mathbf{T}_{\ell,\mathbb{C}}$ is independent of ℓ. In particular, the rank of $\mathbf{G}_\ell$ is independent of ℓ.*

Proof. For any ℓ, ℓ', we can find some $\bar{v}$ such that both $\mathbf{T}_{\bar{v},\ell}$ and $\mathbf{T}_{\bar{v},\ell'}$ are maximal Frobenius tori of $\mathbf{G}_\ell$ and $\mathbf{G}_{\ell'}$ respectively by Corollary 2.2 and Theorem 2.1. By weak compatibility, the tautological representations of $\mathbf{T}_{\bar{v},\ell}(\mathbb{C})$ and $\mathbf{T}_{\bar{v},\ell'}(\mathbb{C})$ are isomorphic. Since all maximal tori of a connected reductive group (over algebraically closed field) are conjugate, we are done. □

Proposition 2.4. [De80],[Se81] *(see also* [LP97]*) The conditions on Frobenius eigenvalues of Theorem 2.1 always hold for E_ℓ-representations arising from geometry.*

2.2. **Locally algebraic abelian representations.** This subsection describes the essence of [Se98, Ch. II, III]. Let $\mathrm{Art} : I_K \to \mathrm{Gal}(\bar{K}/K)^{\mathrm{ab}}$ be the *Artin map* and $i : (K \otimes_{\mathbb{Q}} \mathbb{Q}_\ell)^* \hookrightarrow I_K$ the embedding. Let E_ℓ be a finite extension of $\mathbb{Q}_\ell$ and $\mathbf{T} := \mathrm{Res}_{K/\mathbb{Q}}\mathbb{G}_m$ the $\mathbb{Q}$-torus given by the *Weil restriction of scalars*. For any $\mathbb{Q}$-algebra A, we have the group isomorphism $\mathbf{T}(A) \cong (K \otimes_{\mathbb{Q}} A)^*$.

Definition 3. [Se98, Ch. III §2] A continuous abelian E_ℓ-representation $\rho_\ell : \mathrm{Gal}(\bar{K}/K)^{\mathrm{ab}} \to \mathrm{GL}_n(E_\ell)$ is said to be *locally algebraic* if there exists an E_ℓ-morphism $r : \mathbf{T} \times_{\mathbb{Q}} E_\ell \to \mathrm{GL}_{n,E_\ell}$ such that

$$\rho_\ell \circ \mathrm{Art} \circ i(x) = r(x^{-1}) \tag{2}$$

for all $x \in (K \otimes_{\mathbb{Q}} \mathbb{Q}_\ell)^*$ close enough to the identity (ℓ-adic topology), where we identify $(K \otimes_{\mathbb{Q}} \mathbb{Q}_\ell)^* \subset (K \otimes_{\mathbb{Q}} E_\ell)^* = \mathbf{T}(E_\ell)$.

Local algebraicity of ρ_ℓ depends only on the local representations of ρ_ℓ above ℓ, i.e., the restrictions of ρ_ℓ to the decomposition groups of $\mathrm{Gal}(\bar{K}/K)$ at the places of $\bar{K}$ above ℓ. This property is invariant under field extensions of E_ℓ (obviously) as well as the restriction of scalars.

Proposition 2.5. *(see* [Ri76, §5]*) A continuous abelian E_ℓ-representation $\rho_\ell : \mathrm{Gal}(\bar{K}/K)^{\mathrm{ab}} \to \mathrm{GL}_n(E_\ell)$ is locally algebraic if and only if it is locally algebraic when viewed as a $\mathbb{Q}_\ell$-representation.*

Let S be a finite subset of finite places of K. For any *modulus* $\mathfrak{m} = (m_v)_{v \in S}$ ($m_v \in \mathbb{N}$) with support $\mathrm{Supp}(\mathfrak{m}) = S$, Serre has constructed [Se98, Ch. II] the *Serre group* $\mathbf{S}_\mathfrak{m}$ and an abelian rational representation with values in $\mathbf{S}_\mathfrak{m}$ for each ℓ:

$$\epsilon_\ell : \mathrm{Gal}(\bar{K}/K)^{\mathrm{ab}} \to \mathbf{S}_\mathfrak{m}(\mathbb{Q}_\ell). \tag{3}$$

The Serre group $\mathbf{S}_\mathfrak{m}$ is a diagonalizable group over $\mathbb{Q}$ satisfying the short exact sequence

$$1 \to \mathbf{T}_\mathfrak{m} \to \mathbf{S}_\mathfrak{m} \to C_\mathfrak{m} \to 1, \tag{4}$$

where $\mathbf{T}_\mathfrak{m}$ is the identity component of $\mathbf{S}_\mathfrak{m}$.

Proposition 2.6. [Se98, Ch. II §2] *We list some properties of ϵ_ℓ and $\mathbf{S}_\mathfrak{m}$.*

(i) The representation ϵ_ℓ is unramified outside S and S_ℓ, the places v of K above ℓ.
(ii) The image of the Frobenii above v belong to $\mathbf{S}_\mathfrak{m}(\mathbb{Q})$ for all $v \notin S \cup S_\ell$ and is dense in $\mathbf{S}_\mathfrak{m}$.
(iii) The representations of $\mathbf{S}_\mathfrak{m}$ are always semisimple.
(iv) The finite group $C_\mathfrak{m}$ is the Galois group of the class field of K corresponding to $\mathfrak{m}$.
(v) The dimension d_K of $\mathbf{S}_\mathfrak{m}$ depends only on K, not on $\mathfrak{m}$.

Definition 4. [Se98, Ch. II §2] A continuous abelian E_ℓ-representation $\rho_\ell : \mathrm{Gal}(\bar{K}/K)^{\mathrm{ab}} \to \mathrm{GL}_n(E_\ell)$ is said to be *associated to some Serre group* $\mathbf{S}_\mathfrak{m}$ if there exists an E_ℓ-morphism $\eta_\ell : \mathbf{S}_{\mathfrak{m},E_\ell} \to \mathrm{GL}_{n,E_\ell}$ such that

$$\rho_\ell = \eta_\ell \circ \epsilon_\ell. \tag{5}$$

Theorem 2.7. [Se98, Ch. III Thm. 1, Thm. 2] *Let $\rho_\ell : \mathrm{Gal}(\bar{K}/K)^{\mathrm{ab}} \to \mathrm{GL}_n(E_\ell)$ be a continuous abelian E_ℓ-representation.*

(i) ρ_ℓ is locally algebraic if and only if it is associated to some Serre group $\mathbf{S}_\mathfrak{m}$.
(ii) If ρ_ℓ is locally algebraic and unramified outside a finite set of places S, then it is associated to some Serre group $\mathbf{S}_\mathfrak{m}$ with $\mathrm{Supp}(\mathfrak{m}) = S$.

Definition 5. [Se98, Ch. I §2] Let $\varphi_\ell : \mathrm{Gal}(\bar{K}/K) \to \mathrm{GL}_n(E_\ell)$ be a continuous E_ℓ-representation and E a number field contained in E_ℓ. Then φ_ℓ is said to be *E-rational* if φ_ℓ is unramified outside a finite subset S of finite places v of K and the characteristic polynomial $P_{v,\varphi_\ell}(t)$ of the Frobenii above v belongs to $E[t]$ for all v outside S. The term $\mathbb{Q}$-rational is abbreviated as *rational*.

For any finite place v of a number field, let p_v be the characteristic of v. For any number field E with a finite place λ, let E_λ be the completion of E with respect to λ. An E_λ-representation is also called a λ-adic representation.

Definition 6. [Se98, Ch. I §2] For each finite place λ of E, let φ_λ be an E-rational λ-adic representation of $\mathrm{Gal}(\bar{K}/K)$. The system $\{\varphi_\lambda\}_\lambda$ is said to be *strictly compatible* if there exists a finite subset S of finite places v of K, called the *exceptional set* of $\{\varphi_\lambda\}_\lambda$, such that:

(i) let $S_\lambda = \{v : \ p_v = p_\lambda\}$, then ρ_λ is unramified if $v \notin S \cup S_\lambda$.
(ii) $P_{v,\varphi_\lambda}(t) = P_{v,\varphi_{\lambda'}}(t)$ if $v \notin S \cup S_\lambda \cup S_{\lambda'}$.

Theorem 2.8. [Se98, Ch. II §3.4] *If a continuous abelian E_ℓ-representation $\rho_\ell : \mathrm{Gal}(\bar{K}/K)^{\mathrm{ab}} \to \mathrm{GL}_n(E_\ell)$ is associated to some Serre group $\mathbf{S}_\mathfrak{m}$, then ρ_ℓ is semisimple and E-rational for some subfield E of the Galois closure of K and the $|C_\mathfrak{m}|$th roots of unity in $\bar{\mathbb{Q}}$, where the finite group $C_\mathfrak{m}$ is defined in (4).*

Corollary 2.9. *If a continuous abelian $E_{\ell'}$-representation $\rho_{\ell'} : \mathrm{Gal}(\bar{K}/K)^{\mathrm{ab}} \to \mathrm{GL}_n(E_{\ell'})$ is associated to some Serre group $\mathbf{S}_\mathfrak{m}$, then there exist a strictly compatible E-rational system $\{\rho_\lambda\}_\lambda$ of semisimple abelian λ-adic representations with the exceptional set $\mathrm{Supp}(\mathfrak{m})$ and a local field isometry $E_{\lambda'} \cong E_{\ell'}$ for some λ' such that*

$$\rho_{\ell'} = \rho_{\lambda'}. \tag{6}$$

Proof. By Theorem 2.8 and [Se98, Ch. II §2.4], $\eta_{\ell'}$ (in Definition 4) is the base change to $E_{\ell'}$ of an E-morphism $\eta : \mathbf{S}_{\mathfrak{m},E} \to \mathrm{GL}_{n,E}$ and we may assume the number field E is dense in $E_{\ell'}$. Then η induces the desired system of λ-adic representations (see e.g. [Se98, Ch. II Thm. 1]). Pick λ' such that $E_{\lambda'}$ is isometric to $E_{\ell'}$, we obtain $\rho_{\ell'} = \rho_{\lambda'}$. □

The following *converse* of Theorem 2.8 (in light of Theorem 2.7) follows from the arguments of Serre in [Se98, Ch. III §3] and a result of transcendental numbers by Waldschmidt [Wa81].

Theorem 2.10. *(see e.g.* [He80, Thm. 2]*) If a continuous semisimple abelian E_ℓ-representation $\rho_\ell : \mathrm{Gal}(\bar{K}/K)^{\mathrm{ab}} \to \mathrm{GL}_n(E_\ell)$ is E-rational, then ρ_ℓ is locally algebraic.*

Definition 7. [Se67, §2] Let L be a finite extension of $\mathbb{Q}_\ell$ and $B_{HT} := \bigoplus_{i\in\mathbb{Z}} \mathbb{C}_\ell(i)$ the period ring, where the $\mathrm{Gal}(\bar{L}/L)$-module $\mathbb{C}_\ell(i)$ is the twist of $\mathbb{C}_\ell$ (the completion of $\bar{L}$) by the ith power of the cyclotomic character. A continuous E_ℓ-representation $\varphi_\ell : \mathrm{Gal}(\bar{L}/L) \to \mathrm{GL}(V_\ell)$ is said to be *of Hodge-Tate type* if the natural $\mathrm{Gal}(\bar{L}/L)$-inclusion

$$B_{HT} \otimes_{\mathbb{Q}_\ell} (B_{HT} \otimes_{\mathbb{Q}_\ell} V_\ell)^{\mathrm{Gal}(\bar{L}/L)} \to B_{HT} \otimes_{\mathbb{Q}_\ell} V_\ell \tag{7}$$

is an isomorphism.

Remark 2.11. *The E_ℓ-representation V_ℓ is viewed as a $\mathbb{Q}_\ell$-representation in (7).*

Theorem 2.12. [Ta67],[Se98, Ch. III Appendix] *A continuous abelian $\mathbb{Q}_\ell$-representation $\rho_\ell : \mathrm{Gal}(\bar{K}/K)^{\mathrm{ab}} \to \mathrm{GL}_n(\mathbb{Q}_\ell)$ is locally algebraic if and only if the local representations of $\varphi_\ell : \mathrm{Gal}(\bar{K}/K) \twoheadrightarrow \mathrm{Gal}(\bar{K}/K)^{\mathrm{ab}} \xrightarrow{\rho_\ell} \mathrm{GL}_n(\mathbb{Q}_\ell)$ above ℓ are*

(i) of Hodge-Tate type and
(ii) semisimple when restricted to the inertia groups.

Given a continuous semisimple abelian E_ℓ-representation

$$\varphi_\ell : \mathrm{Gal}(\bar{K}/K) \twoheadrightarrow \mathrm{Gal}(\bar{K}/K)^{\mathrm{ab}} \xrightarrow{\rho_\ell} \mathrm{GL}_n(E_\ell), \tag{8}$$

the results of this subsection are summarized by the diagram.

$$\begin{array}{ccccc} \mathrm{B} & \overset{2.5}{\Longleftrightarrow} & \mathrm{A} & \overset{2.10}{\Longleftarrow} & \mathrm{D} \\ 2.12 \Updownarrow & & 2.7(i) \Updownarrow & \nearrow_{2.8} & \Uparrow \text{obvious} \\ \mathrm{F} & & \mathrm{C} & \underset{2.9}{\Longrightarrow} & \mathrm{E} \end{array} \tag{9}$$

A. ρ_ℓ is locally algebraic;
B. ρ_ℓ is locally algebraic as a $\mathbb{Q}_\ell$-representation;
C. ρ_ℓ is associated to some Serre group $\mathbf{S}_{\mathfrak{m}}$;
D. ρ_ℓ is E-rational;
E. ρ_ℓ can be extended to a strictly compatible E-rational system of semisimple abelian λ-adic representations with the exceptional set $\mathrm{Supp}(\mathfrak{m})$;
F. The local representations above ℓ of φ_ℓ are of Hodge-Tate type.

2.3. **Formal bi-character.** Let $\mathbf{G}_{\mathbb{C}}$ be a connected reductive Lie subgroup of $\mathrm{GL}_n(\mathbb{C})$ and $\mathbf{G}_{\mathbb{C}}^{\mathrm{der}}$ the derived group (semisimple part) of $\mathbf{G}_{\mathbb{C}}$. Let $\mathbf{T}_{\mathbb{C}}$ be a maximal torus of $\mathbf{G}_{\mathbb{C}}$ and $\mathbf{T}_{\mathbb{C}}^{\mathrm{ss}} := (\mathbf{T}_{\mathbb{C}} \cap \mathbf{G}_{\mathbb{C}}^{\mathrm{der}})^\circ$ the maximal torus of $\mathbf{G}_{\mathbb{C}}^{\mathrm{der}}$. Since all maximal tori of $\mathbf{G}_{\mathbb{C}}$ are conjugate, the following definition makes sense.

Definition 8. [Hui15b, Defs. 5,6]

(i) The *formal character* of $\mathbf{G}_{\mathbb{C}} \hookrightarrow \mathrm{GL}_n(\mathbb{C})$ is defined as the tautological representation $\mathbf{T}_{\mathbb{C}} \hookrightarrow \mathrm{GL}_n(\mathbb{C})$.
(ii) The *formal bi-character* of $\mathbf{G}_{\mathbb{C}} \hookrightarrow \mathrm{GL}_n(\mathbb{C})$ is defined as the tautological representation $\mathbf{T}_{\mathbb{C}}^{\mathrm{ss}} \subset \mathbf{T}_{\mathbb{C}} \hookrightarrow \mathrm{GL}_n(\mathbb{C})$.

Let E_ℓ be a finite extension of $\mathbb{Q}_\ell$ and fix an embedding $E_\ell \hookrightarrow \mathbb{C}$ for all ℓ. Consider a weakly compatible system of semisimple E_ℓ-representations

$$\psi_\ell : \mathrm{Gal}(\bar{K}/K) \to \mathrm{GL}_n(E_\ell), \quad \forall \ell \tag{10}$$

and denote by $\mathbf{G}_\ell$ the algebraic monodromy group of ψ_ℓ.

Theorem 2.13. *Suppose $\{\psi_\ell\}_\ell$ satisfies the following conditions:*

(i) $\mathbf{G}_\ell$ is connected for all ℓ;
(ii) for each ℓ, the eigenvalues of $\psi_\ell(\mathrm{Frob}_{\bar{v}})$ satisfy the conditions in Theorem 2.1 for almost all finite places v of K;
(iii) the local representations above ℓ of ψ_ℓ are of Hodge-Tate type for all ℓ.

Then the formal bi-character of $\mathbf{G}_\ell(\mathbb{C}) \hookrightarrow \mathrm{GL}_n(\mathbb{C})$ *is independent of* ℓ.

Remark 2.14. *In case* $\{\psi_\ell\}_\ell$ *is only a strictly compatible rational system of* $\mathbb{Q}_\ell$*-representations without the three conditions, Theorem 2.13 follows from* [Hui13, Thm. 3.19]. *The added conditions allow us to streamline the proof of* [Hui13, Thm. 3.19]*, in which we follow closely for Theorem 2.13.*

Proof. Identify $\mathbf{G}_\ell$ as a connected reductive subgroup of GL_{n,E_ℓ} by (i) and the semisimplicity. Then the derived group $\mathbf{G}_\ell^{\mathrm{der}}$ is semisimple. Let $\eta : \mathbf{S}_{\mathfrak{m}} \to \mathrm{GL}_{m,\mathbb{Q}}$ be a faithful representation and form a strictly compatible system of semisimple abelian E_ℓ-representations $\{\rho_\ell := \eta_\ell \circ \epsilon_\ell\}_\ell$ (Definition 4). Then $\{\rho_\ell\}_\ell$ satisfies (ii) by [Se98, Ch. II §3.4 Prop. 2] and (iii) by the diagram (9), so does the weakly compatible system $\{\psi_\ell \oplus \rho_\ell\}_\ell$. Since $\psi_\ell \oplus \rho_\ell$ is semisimple, the algebraic monodromy group $\mathbf{H}_\ell$ is reductive. Since ρ_ℓ is abelian, the derived group $\mathbf{H}_\ell^{\mathrm{der}}$ of $\mathbf{H}_\ell$ is exactly equal to $\mathbf{G}_\ell^{\mathrm{der}}$ (or $\mathbf{G}_\ell^{\mathrm{der}} \times \{1\} \subset \mathrm{GL}_{n,E_\ell} \times \mathrm{GL}_{m,E_\ell}$). Since $\mathbf{H}_\ell$ is reductive, there exist $k \in \mathbb{N}$ and a subrepresentation W_ℓ of $(\psi_\ell \oplus \rho_\ell)^{\otimes k} = (E_\ell^{n+m})^{\otimes k}$ such that

$$\mathbf{G}_\ell^{\mathrm{der}} = \ker(\mathbf{H}_\ell \to \mathrm{GL}_{W_\ell})^\circ. \tag{11}$$

Since W_ℓ is abelian semisimple and its local representations above ℓ are of Hodge-Tate type by construction, W_ℓ is associated to some Serre group by the diagram (9), which implies that the center $Z(\mathbf{H}_\ell^\circ)$ of the identity component $\mathbf{H}_\ell^\circ$ satisfies the inequality

$$\dim Z(\mathbf{H}_\ell^\circ) \leq \dim \mathbf{S}_{\mathfrak{m},E_\ell} \overset{2.6(v)}{=:} d_K. \tag{12}$$

Since the projection of $\mathbf{H}_\ell \subset \mathrm{GL}_{n,E_\ell} \times \mathrm{GL}_{m,E_\ell}$ to the second factor is $\mathbf{S}_{\mathfrak{m},E_\ell}$ by Proposition 2.6(ii), we conclude with (12) that

$$\dim Z(\mathbf{H}_\ell^\circ) = d_K. \tag{13}$$

Since $\{\psi_\ell \oplus \rho_\ell\}_\ell$ satisfies (ii) (the Frobenius eigenvalues conditions), the restriction $\{(\psi_\ell \oplus \rho_\ell)|_{\mathrm{Gal}(\bar{K}/L)}\}_\ell$ satisfies (ii) and weak compatibility for any finite extension L of K. By Corollary 2.2, we can find some finite extension L such that the algebraic monodromy group of $(\psi_\ell \oplus \rho_\ell)|_{\mathrm{Gal}(\bar{K}/L)}$ is the identity component $\mathbf{H}_\ell^\circ$ for all ℓ. By applying Corollary 2.3 to the system $\{(\psi_\ell \oplus \rho_\ell)|_{\mathrm{Gal}(\bar{K}/L)}\}_\ell$, we obtain a maximal torus $\mathbf{T}_{\ell,\mathbb{C}}$ (Frobenius torus) of the connected reductive Lie group $\mathbf{H}_\ell^\circ(\mathbb{C})$, in which the construction implies that the tautological representation

$$\mathbf{T}_{\ell,\mathbb{C}} \hookrightarrow \mathrm{GL}_n(\mathbb{C}) \times \mathrm{GL}_m(\mathbb{C}) \tag{14}$$

is independent of ℓ, i.e., for all ℓ', ℓ'', $\mathbf{T}_{\ell',\mathbb{C}}$ and $\mathbf{T}_{\ell'',\mathbb{C}}$ are conjugate in $\mathrm{GL}_n(\mathbb{C}) \times \mathrm{GL}_m(\mathbb{C})$. If we let π_1 (resp. π_2) be the projection of $\mathrm{GL}_n(\mathbb{C}) \times \mathrm{GL}_m(\mathbb{C})$ to the first (resp. second) factor, then the ℓ-independence of (14) implies the tautological representation

$$\pi_1(\ker(\pi_2|_{\mathbf{T}_{\ell,\mathbb{C}}})^\circ) \subset \pi_1(\mathbf{T}_{\ell,\mathbb{C}}) \hookrightarrow \mathrm{GL}_n(\mathbb{C}) \tag{15}$$

is independent of ℓ.

Since $\pi_1(\mathbf{H}_\ell) = \mathbf{G}_\ell$, $\pi_1(\mathbf{T}_{\ell,\mathbb{C}})$ is a maximal torus of $\mathbf{G}_\ell(\mathbb{C})$. Since we have $\pi_2(\mathbf{H}_\ell) = \mathbf{S}_{\mathfrak{m},E_\ell}$ and (13), we deduce that

$$\ker(\pi_2 : \mathbf{H}_\ell \to \mathrm{GL}_{m,E_\ell})^\circ = \mathbf{H}_\ell^{\mathrm{der}} = \mathbf{G}_\ell^{\mathrm{der}} \times \{1\}. \tag{16}$$

Hence, $\pi_1(\ker(\pi_2|_{\mathbf{T}_{\ell,\mathbb{C}}})^\circ)$ is a maximal torus of $\mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C})$. Therefore, we conclude that (15), the formal bi-character of $\mathbf{G}_\ell(\mathbb{C}) \hookrightarrow \mathrm{GL}_n(\mathbb{C})$, is independent of ℓ. □

Suppose $\mathbf{G}_\mathbb{C}$ and $\mathbf{G}'_\mathbb{C}$ are two connected reductive Lie subgroups of $\mathrm{GL}_n(\mathbb{C})$ such that their formal bi-characters are isomorphic. After conjugating $\mathbf{G}'_\mathbb{C}$ by an element of $\mathrm{GL}_n(\mathbb{C})$, we may assume

$$\mathbf{T}_\mathbb{C}^{\mathrm{ss}} \subset \mathbf{T}_\mathbb{C} \hookrightarrow \mathrm{GL}_n(\mathbb{C}) \tag{17}$$

is the common formal bi-character of $\mathbf{G}_\mathbb{C}$ and $\mathbf{G}'_\mathbb{C}$, meaning that $\mathbf{T}_\mathbb{C}$ (resp. $\mathbf{T}_\mathbb{C}^{\mathrm{ss}}$) is a common maximal torus of $\mathbf{G}_\mathbb{C}$ and $\mathbf{G}'_\mathbb{C}$ (resp. $\mathbf{G}_\mathbb{C}^{\mathrm{der}}$ and $\mathbf{G}_\mathbb{C}^{\prime\mathrm{der}}$). Let Γ be the character group of $\mathbf{T}_\mathbb{C}^{\mathrm{ss}}$ and $R \subset \Gamma$ (resp. $R' \subset \Gamma$) the roots of $\mathbf{G}_\mathbb{C}^{\mathrm{der}}$ (resp. $\mathbf{G}_\mathbb{C}^{\prime\mathrm{der}}$) with respect to $\mathbf{T}_\mathbb{C}^{\mathrm{ss}}$.

Proposition 2.15. [Hui15b, Cor. 3.8] *Let $\mathbf{G}_\mathbb{C}$ and $\mathbf{G}'_\mathbb{C}$ be as above. If $R = R'$ in Γ, then $\mathbf{G}_\mathbb{C}$ and $\mathbf{G}'_\mathbb{C}$ are conjugate in $\mathrm{GL}_n(\mathbb{C})$.*

We give sufficient conditions for $R = R'$ in Γ.

Proposition 2.16. *Let $\mathbf{G}_\mathbb{C}$ and $\mathbf{G}'_\mathbb{C}$ be as above. Suppose the following conditions hold:*

(i) $\mathbf{G}_\mathbb{C}^{\mathrm{der}}$ and $\mathbf{G}_\mathbb{C}^{\prime\mathrm{der}}$ are conjugate in $\mathrm{GL}_n(\mathbb{C})$;
(ii) the roots R of $\mathbf{G}_\mathbb{C}^{\mathrm{der}}$ are stable under the normalizer $N_{\mathrm{GL}_n(\mathbb{C})}(\mathbf{T}_\mathbb{C}^{\mathrm{ss}})$.

Then $R = R'$ in Γ.

Proof. Since $\mathbf{G}_\mathbb{C}^{\mathrm{der}}$ and $\mathbf{G}_\mathbb{C}^{\prime\mathrm{der}}$ are conjugate in $\mathrm{GL}_n(\mathbb{C})$ by (i), they are conjugate by an element of $N_{\mathrm{GL}_n(\mathbb{C})}(\mathbf{T}_\mathbb{C}^{\mathrm{ss}})$, inducing an automorphism of Γ mapping R to R'. Hence, $R = R'$ by (ii). □

Back to the semisimple weakly compatible system (10): $\{\psi_\ell : \mathrm{Gal}(\bar{K}/K) \to \mathrm{GL}_n(E_\ell)\}_\ell$. Let $\mathbf{T}_\mathbb{C}^{\mathrm{ss}}$ be a maximal torus of $\mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C})$ and R the roots of $\mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C})$ with respect to $\mathbf{T}_\mathbb{C}^{\mathrm{ss}}$.

Corollary 2.17. *Suppose $\{\psi_\ell\}_\ell$ satisfies the following conditions:*

(i) $\mathbf{G}_\ell$ is connected for all ℓ;
(ii) for each ℓ, the eigenvalues of $\psi_\ell(\mathrm{Frob}_v)$ satisfy the conditions in Theorem 2.1 for almost all finite places v of K;
(iii) the local representations above ℓ of ψ_ℓ are of Hodge-Tate type for all ℓ;
(iv) the conjugacy class of $\mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C})$ in $\mathrm{GL}_n(\mathbb{C})$ is independent of ℓ;
(v) the roots R of $\mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C})$ are stable under the normalizer $N_{\mathrm{GL}_n(\mathbb{C})}(\mathbf{T}_\mathbb{C}^{\mathrm{ss}})$.

Then the conjugacy class of $\mathbf{G}_\ell(\mathbb{C})$ in $\mathrm{GL}_n(\mathbb{C})$ is independent of ℓ. In particular, the dimension of the abelian part $(E_\ell^n)^{\mathrm{ab}}$ (resp. the invariants $(E_\ell^n)^{\mathrm{Gal}(\bar{K}/K)}$) is independent of ℓ.

Proof. The formal bi-character of $\mathbf{G}_\ell(\mathbb{C}) \hookrightarrow \mathrm{GL}_n(\mathbb{C})$ is independent of ℓ by (i),(ii),(iii) and Theorem 2.13. Then (iv),(v) and Propositions 2.16, 2.15 imply that the conjugacy class of $\mathbf{G}_\ell(\mathbb{C})$ in $\mathrm{GL}_n(\mathbb{C})$ is independent of ℓ. □

2.4. **Proofs of the theorems.**

Theorem 1.1. Let $\{V_\ell\}_\ell$ be a strictly compatible rational system arising from geometry, with the exceptional set S. Let $\mathbf{G}_\ell$ be the algebraic monodromy group of V_ℓ^{ss} for all ℓ. Suppose the following conditions hold:

(i) $\mathbf{G}_\ell$ is connected for all ℓ;

(ii) the conjugacy class of $\mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C})$ in $\mathrm{GL}_N(\mathbb{C})$ is independent of ℓ;
(iii) the roots R of $\mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C})$ are stable under the normalizer $N_{\mathrm{GL}_N(\mathbb{C})}(\mathbf{T}_{\mathbb{C}}^{\mathrm{ss}})$.

Then the abelian part $\{V_\ell^{\mathrm{ab}}\}_\ell$ is strictly compatible rational with the same exceptional set S.

Proof. Since $\{V_\ell\}_\ell$ is strictly compatible rational arising from geometry, $\{V_\ell^{\mathrm{ss}}\}_\ell$ satisfies 2.17(ii) and 2.17(iii) respectively by Proposition 2.4 and the Hodge-Tate conjecture proved by Faltings [Fa88]. Our conditions (i),(ii),(iii) are just 2.17(i),(iv),(v). Hence, $\dim V_\ell^{\mathrm{ab}}$ is independent of ℓ by Corollary 2.17. Let V_ℓ^{c} be the complementary representation of V_ℓ^{ab} in V_ℓ^{ss}. Let χ_ℓ be a one dimensional $\bar{\mathbb{Q}}_\ell$-representation (character) of $\mathrm{Gal}(\bar{K}/K)$. By the construction of V_ℓ^{c}, the twisted representation

$$V_\ell^{\mathrm{c}} \otimes_{\mathbb{Q}_\ell} \chi_\ell \text{ cannot have one dimensional subrepresentation.} \tag{18}$$

Pick a prime ℓ' not dividing any place in S and a finite extension $E_{\ell'}$ of $\mathbb{Q}_{\ell'}$ such that $V_{\ell'}^{\mathrm{ss}} \otimes_{\mathbb{Q}_{\ell'}} E_{\ell'}$ decomposes as a direct sum of absolutely irreducible representations. Consider the isomorphism classes $\mathcal{X}_{\ell'}$ of the characters of $V_{\ell'}^{\mathrm{ab}} \otimes_{\mathbb{Q}_{\ell'}} E_{\ell'}$. For any $W_{\ell'} \in \mathcal{X}_{\ell'}$, $W_{\ell'}$ is unramified outside $S \cup S_{\ell'}$ (Definition 6(i)). Since $W_{\ell'}$ arises from geometry, its local representations above ℓ' are of Hodge-Tate type [Fa88]. By the diagram (9) and Theorem 2.7(ii), $W_{\ell'}$ is associated to some Serre group $\mathbf{S}_{\mathfrak{m}}$ with $\mathrm{Supp}(\mathfrak{m}) = S \cup S_{\ell'}$. Since $\mathcal{X}_{\ell'}$ is finite, there exists a number field E dense in $E_{\ell'}$ such that for every $W_{\ell'} \in \mathcal{X}_{\ell'}$, $W_{\ell'}$ extends to a strictly compatible E-rational system of semisimple abelian λ-adic representations $\{W_\lambda\}_\lambda$ with the exceptional set $S \cup S_{\ell'}$ by the diagram (9). Suppose $\{W_{1,\lambda}\}_\lambda, \{W_{2,\lambda}\}_\lambda$ are the λ-adic systems corresponding to $W_{1,\ell'}, W_{2,\ell'} \in \mathcal{X}_{\ell'}$. By the strict compatibility of the systems,

$$\text{if } W_{1,\ell'} \ncong W_{2,\ell'}, \text{ then } W_{1,\lambda} \ncong W_{2,\lambda} \text{ for all } \lambda. \tag{19}$$

Let λ' be the prime of E corresponding to $E \hookrightarrow E_{\ell'}$. For every prime ℓ other than ℓ', fix a λ that divides ℓ and let E_ℓ be the completion of E with respect to λ. Then we obtain for every $W_{\ell'} \in \mathcal{X}_{\ell'}$, a subfamily $\{W_\ell\}_\ell$ of $\{W_\lambda\}_\lambda$, indexed by ℓ, satisfying the following properties.

Proposition 2.18. *The system $\{W_\ell\}_\ell$ extends $W_{\ell'}$ and is a weakly compatible system of E_ℓ-representations. Moreover, W_ℓ is unramified outside $(S \cup S_{\ell'}) \cup S_\ell$.*

The invariant dimension of the representation $\mathrm{Hom}_{E_\ell}(W_\ell, V_\ell^{\mathrm{ss}} \otimes_{\mathbb{Q}_\ell} E_\ell) = V_\ell^{\mathrm{ss}} \otimes_{\mathbb{Q}_\ell} W_\ell^*$, that is,

$$\dim_{E_\ell}(V_\ell^{\mathrm{ss}} \otimes_{\mathbb{Q}_\ell} W_\ell^*)^{\mathrm{Gal}(\bar{K}/K)}, \tag{20}$$

counts the number of copies of W_ℓ in V_ℓ^{ab} by (18). Assume (20) is independent of ℓ for each $W_{\ell'} \in \mathcal{X}_{\ell'}$, then the facts that $\{W_\ell\}_\ell$ extends $W_{\ell'}$ and $\dim V_\ell^{\mathrm{ab}}$ is independent of ℓ, together with (19) and Proposition 2.18, imply that for any pair of primes ℓ'', ℓ''',

$$P_{v,V_{\ell''}^{\mathrm{ab}}}(t) = P_{v,V_{\ell'''}^{\mathrm{ab}}}(t) \in E[t] \text{ if } v \notin (S \cup S_{\ell'}) \cup S_{\ell''} \cup S_{\ell'''}. \tag{21}$$

Roughly speaking, it says that the eigenvalues of Frobenius on V_ℓ^{ab} is independent of ℓ. Let $P_v(t)$ be the characteristic polynomial in (21). Then $P_v(t) \in \mathbb{Q}_\ell[t]$ for almost all ℓ because V_ℓ^{ab} is a $\mathbb{Q}_\ell$-vector space. Since $P_v(t) \in E[t]$ and E is a finite extension of $\mathbb{Q}$, we obtain $P_v(t) \in \mathbb{Q}[t]$. Thus, $\{V_\ell^{\mathrm{ab}}\}_\ell$ is a strictly compatible rational with the exceptional set $S \cup S_{\ell'}$. By picking another prime ℓ'' not equal to ℓ' in the second paragraph of this proof and repeat

the above arguments, we conclude that $\{V_\ell^{\mathrm{ab}}\}_\ell$ is a strictly compatible rational with the exceptional set S.

To complete the proof, it remains to prove that (20) is independent of ℓ for each $W_{\ell'} \in \mathcal{X}_{\ell'}$. It suffices to verify the five conditions of Corollary 2.17 for the semisimple weakly compatible system $\{V_\ell^{\mathrm{ss}} \otimes_{\mathbb{Q}_\ell} W_\ell^*\}_\ell$ for any $W_{\ell'} \in \mathcal{X}_{\ell'}$. Let $\mathbf{H}_\ell$ be the algebraic monodromy group of $V_\ell^{\mathrm{ss}} \otimes_{\mathbb{Q}_\ell} W_\ell^*$. The natural morphism $V_\ell^{\mathrm{ss}} \otimes_{\mathbb{Q}_\ell} E_\ell \to V_\ell^{\mathrm{ss}} \otimes_{\mathbb{Q}_\ell} W_\ell^*$ implies that $\mathbf{G}_\ell$ surjects onto $\mathbf{H}_\ell$. Since $\mathbf{G}_\ell$ is connected by (i), $\mathbf{H}_\ell$ is also connected and 2.17(i) holds. Since $V_\ell^{\mathrm{ss}} \otimes_{\mathbb{Q}_\ell} W_\ell^*$ and $V_\ell^{\mathrm{ss}} \otimes_{\mathbb{Q}_\ell} E_\ell$ are differed by a twist of the character W_ℓ^*, the derived subgroups of the Galois images are equal under suitable identification, which implies $\mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C}) = \mathbf{H}_\ell^{\mathrm{der}}(\mathbb{C})$ in $\mathrm{GL}_N(\mathbb{C})$. The conditions 2.17(iv),(v) then follow from our (ii),(iii). Since $\{W_\ell\}_\ell$ arises from an E-representation of some Serre group $\mathbf{S}_{\mathfrak{m},E}$, $\{W_\ell\}_\ell$ satisfies 2.17(ii) by [Se98, Ch. II §3.4] and 2.17(iii) by the diagram (9). Hence, $V_\ell^{\mathrm{ss}} \otimes_{\mathbb{Q}_\ell} W_\ell^*$ also satisfies 2.17(ii),(iii). We conclude by Corollary 2.17 that the invariant dimension (20) is independent of ℓ for each $W_{\ell'} \in \mathcal{X}_{\ell'}$. □

Theorem 2.19. [Hui15b, §3] *Let $\mathbf{G}_{\mathbb{C}}$ and $\mathbf{G}'_{\mathbb{C}}$ be two connected reductive Lie subgroup of $\mathrm{GL}_N(\mathbb{C})$ having the common formal bi-character $\mathbf{T}_{\mathbb{C}}^{\mathrm{ss}} \subset \mathbf{T}_{\mathbb{C}} \hookrightarrow \mathrm{GL}_N(\mathbb{C})$. Suppose $\mathbf{G}_{\mathbb{C}}'^{\mathrm{der}}$ satisfies the Lie type conditions in Hypothesis A, i.e.,*

- *$\mathbf{G}_{\mathbb{C}}'^{\mathrm{der}}$ has at most one factor of type A_4;*
- *if $\mathbf{H}_{\mathbb{C}}$ is an almost simple factor of $\mathbf{G}_{\mathbb{C}}'^{\mathrm{der}}$, then $\mathbf{H}_{\mathbb{C}}$ is of type A_n for some $n \in \mathbb{N}\backslash\{1,2,3,5,7,8\}$.*

Then $\mathbf{G}_{\mathbb{C}}$ and $\mathbf{G}'_{\mathbb{C}}$ are conjugate in $\mathrm{GL}_N(\mathbb{C})$ and their roots with respect to $\mathbf{T}_{\mathbb{C}}$ are identical. In particular, $\mathbf{G}_{\mathbb{C}}^{\mathrm{der}}$ and $\mathbf{G}_{\mathbb{C}}'^{\mathrm{der}}$ are conjugate in $\mathrm{GL}_N(\mathbb{C})$ and their roots with respect to $\mathbf{T}_{\mathbb{C}}^{\mathrm{ss}}$ are identical.

Theorem 1.3. Let $\{V_\ell\}_\ell$ be a strictly compatible rational system arising from geometry, with the exceptional set S. Let $\mathbf{G}_\ell$ be the algebraic monodromy group of V_ℓ^{ss} for all ℓ. Suppose (i) $\mathbf{G}_\ell$ is connected for all ℓ and one of the following conditions hold:

(ii) Hypothesis A: there exists some ℓ' such that $\mathbf{G}_{\ell'}^{\mathrm{der}}(\mathbb{C})$ has at most one factor of type A_4 and if $\mathbf{H}_{\mathbb{C}}$ is an almost simple factor of $\mathbf{G}_{\ell'}^{\mathrm{der}}(\mathbb{C})$, then $\mathbf{H}_{\mathbb{C}}$ is of type A_n for some $n \in \mathbb{N}\backslash\{1,2,3,5,7,8\}$;

(ii') the conjugacy class of $\mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C})$ in $\mathrm{GL}_N(\mathbb{C})$ is independent of ℓ and $\mathbf{G}_\ell^{\mathrm{der}}(\mathbb{C})$ is almost simple of type different from A_7, A_8, B_4, D_8.

Then the abelian part $\{V_\ell^{\mathrm{ab}}\}_\ell$ is strictly compatible rational with the same exceptional set S.

Proof. (i+ii): It suffices to show that Hypothesis A implies Theorem 1.1(ii),(iii). This is true by Theorem 2.13 and Theorem 2.19.

(i+ii'): It follows directly from Theorems 1.1 and 3.2. □

Theorem 1.5. Let $\{V_\ell\}_\ell$ be a strictly compatible rational system arising from geometry. Let $\mathbf{G}_\ell$ be the algebraic monodromy group of V_ℓ^{ss} for all ℓ. Suppose (i) $\mathbf{G}_\ell$ is connected for all ℓ and (ii) Hypothesis A is satisfied. Let $\bigoplus^m \bar{W}_{\ell'}$ be a $\bar{\mathbb{Q}}_{\ell'}$-subrepresentation of $V_{\ell'}^{\mathrm{ss}} \otimes \bar{\mathbb{Q}}_{\ell'}$ for some ℓ' and $\{\bar{W}_\ell\}_\ell$ a weakly compatible system of irreducible $\bar{\mathbb{Q}}_\ell$-representations satisfying the following conditions:

(iii) $\{\bar{W}_\ell\}_\ell$ extends $\bar{W}_{\ell'}$;

(iv) for all ℓ, descend $\bar{W}_\ell$ to an E_ℓ-representation W_ℓ in which E_ℓ is a finite extension of $\mathbb{Q}_\ell$. Then the local representations of W_ℓ above ℓ are of Hodge-Tate type.

Then $\{\bigoplus^m \bar{W}_\ell\}_\ell$ is a subsystem of $\{V_\ell^{\mathrm{ss}} \otimes \bar{\mathbb{Q}}_\ell\}_\ell$.

Proof. The idea is similar to Theorem 1.1. Consider the semisimple weakly compatible system

$$\{\mathrm{Hom}_{E_\ell}(W_\ell, V_\ell^{\mathrm{ss}}) = V_\ell^{\mathrm{ss}} \otimes_{\mathbb{Q}_\ell} W_\ell^*\}_\ell. \tag{22}$$

Let $\mathbf{H}_\ell$ be the algebraic monodromy group of $V_\ell^{\mathrm{ss}} \otimes_{\mathbb{Q}_\ell} W_\ell^*$. As $\{W_\ell\}_\ell$ is absolutely irreducible extending $W_{\ell'}$, it suffices to show that invariant dimension

$$\dim_{E_\ell}(V_\ell^{\mathrm{ss}} \otimes_{\mathbb{Q}_\ell} W_\ell^*)^{\mathrm{Gal}(\bar{K}/K)} \tag{23}$$

is independent of ℓ, which follows from Corollary 2.17 once we check the five conditions for the system (22).

The natural morphism $V_\ell^{\mathrm{ss}} \otimes_{\mathbb{Q}_\ell} E_\ell \to V_\ell^{\mathrm{ss}} \otimes_{\mathbb{Q}_\ell} W_\ell^*$ induces the surjection $\mathbf{G}_\ell \to \mathbf{H}_\ell$. Since $\mathbf{G}_\ell$ is connected for all ℓ (i), $\mathbf{H}_\ell$ is also connected and we obtain 2.17(i). Since $W_{\ell'}$ is arises from geometry and $\{W_\ell\}_\ell$ is weakly compatible satisfying (iv), the system (22) satisfies 2.17(ii),(iii). Theorem 2.13 then implies that

$$\text{the formal bi-character of } \mathbf{H}_\ell(\mathbb{C}) \hookrightarrow \mathrm{GL}_M(\mathbb{C}) \text{ is independent of } \ell \tag{24}$$

if M is the dimension of (22). Since $\{\mathbf{G}_\ell\}_\ell$ satisfies Hypothesis A and $\mathbf{G}_\ell$ surjects onto $\mathbf{H}_\ell$ for all ℓ, the family $\{\mathbf{H}_\ell(\mathbb{C})\}_\ell$ of connected reductive Lie groups also satisfies Hypothesis A. Therefore, 2.17(iv),(v) follow from Theorem 2.19 and (24). □

3. Geometry of roots

Let $\mathbf{G}_\mathbb{C}$ be a connected semisimple complex Lie subgroup of $\mathrm{GL}_N(\mathbb{C})$ and $\mathbf{T}_\mathbb{C}$ a maximal torus of $\mathbf{G}_\mathbb{C}$. Denote by Γ the character group of $\mathbf{T}_\mathbb{C}$ and by $R \subset \Gamma$ the roots of $\mathbf{G}_\mathbb{C}$ with respect to $\mathbf{T}_\mathbb{C}$. Let $f : \mathbf{G}_\mathbb{C} \hookrightarrow \mathrm{GL}_N(\mathbb{C})$ be the tautological representation. Then the normalizer $N_{\mathrm{GL}_N(\mathbb{C})}(\mathbf{T}_\mathbb{C})$ acts on Γ. We are interested in the following question:

Q: For what pair $(\mathbf{G}_\mathbb{C}, f)$ is R stable under $N_{\mathrm{GL}_N(\mathbb{C})}(\mathbf{T}_\mathbb{C})$?

Theorem 3.1 (a consequence of Theorem 2.19) below answers **Q** for some type A semisimple $\mathbf{G}_\mathbb{C}$. The goal of this section is to study **Q** under the assumption that $\mathbf{G}_\mathbb{C}$ is almost simple (Theorem 3.2), i.e., the Lie algebra $\mathfrak{g}$ of $\mathbf{G}_\mathbb{C}$ is a complex simple Lie algebra. In the Cartan-Killing classification, $\mathfrak{g}$ is one of A_n ($n \geq 1$), B_n ($n \geq 2$), C_n ($n \geq 3$), D_n ($n \geq 4$), E_6, E_7, E_8, F_4, and G_2.

Theorem 3.1. *Let $\mathbf{G}_\mathbb{C}$ be a connected semisimple complex Lie group satisfying the Lie type conditions in Hypothesis A, i.e.,*

- *$\mathbf{G}_\mathbb{C}$ has at most one factor of type A_4;*
- *if $\mathbf{H}_\mathbb{C}$ is an almost simple factor of $\mathbf{G}_\mathbb{C}$, then $\mathbf{H}_\mathbb{C}$ is of type A_n for some $n \in \mathbb{N}\backslash\{1, 2, 3, 5, 7, 8\}$.*

Then R is stable under $N_{\mathrm{GL}_N(\mathbb{C})}(\mathbf{T}_\mathbb{C})$ for all embedding f of $\mathbf{G}_\mathbb{C}$.

Theorem 3.2. *Let $\mathbf{G}_{\mathbb{C}}$ be a connected almost simple complex Lie group.*

(i) If the type of $\mathbf{G}_{\mathbb{C}}$ is different from A_7, A_8, B_4, D_8, then R is stable under $N_{\mathrm{GL}_N(\mathbb{C})}(\mathbf{T}_{\mathbb{C}})$ for all embeddings f of $\mathbf{G}_{\mathbb{C}}$.

(ii) For all $\mathfrak{g} \in \{A_7, A_8, B_4, D_8\}$, there exists a pair $(\mathbf{G}_{\mathbb{C}}, f)$ such that $\mathbf{G}_{\mathbb{C}}$ is of type $\mathfrak{g}$ and R is not stable under $N_{\mathrm{GL}_N(\mathbb{C})}(\mathbf{T}_{\mathbb{C}})$.

3.1. **A geometric viewpoint.** We introduce a geometric viewpoint on the configuration of R in $\Gamma \otimes_{\mathbb{Z}} \mathbb{R}$ that we learned from [LP90, §1] and developed in [Hui13, §2]. This viewpoint is also the starting point of Theorem 2.19, see [Hui13, §2],[Hui15b, §3]. Let Γ be the character group of the maximal torus $\mathbf{T}_{\mathbb{C}}$, $\mathbb{Z}[\Gamma]$ the group ring, and R (resp. W) the roots (resp. the Weyl group) of the connected complex semisimple $\mathbf{G}_{\mathbb{C}}$ with respect to $\mathbf{T}_{\mathbb{C}}$. Then the formal character $\mathbf{T}_{\mathbb{C}} \hookrightarrow \mathrm{GL}_N(\mathbb{C})$ of $f : \mathbf{G}_{\mathbb{C}} \hookrightarrow \mathrm{GL}_N(\mathbb{C})$ (Definition 8) is equivalent to the element

$$\mathrm{Char}(f) := \alpha_1 + \cdots + \alpha_N \in \mathbb{Z}[\Gamma], \tag{25}$$

where $\alpha_1, ..., \alpha_N$ are the multiset of the weights of f with respect to $\mathbf{T}_{\mathbb{C}}$. Consider the $\mathbb{R}$-vector space $\Gamma \otimes_{\mathbb{Z}} \mathbb{R}$. The formal character defines a positive definite inner product $(\ ,\ )$ on the dual space $(\Gamma \otimes_{\mathbb{Z}} \mathbb{R})^*$ by

$$(x_1^*, x_2^*) = \sum_{i=1}^{N} \alpha_i(x_1^*)\alpha_i(x_2^*). \tag{26}$$

This non-degenerate pairing induces an isomorphism $\Gamma \otimes_{\mathbb{Z}} \mathbb{R} \to (\Gamma \otimes_{\mathbb{Z}} \mathbb{R})^*$ mapping x to x^*. Then we obtain a positive definite inner product $\langle\ ,\ \rangle$ on $\Gamma \otimes_{\mathbb{Z}} \mathbb{R}$ such that $\langle x_1, x_2\rangle = (x_1^*, x_2^*)$ for all $x_1, x_2 \in \Gamma \otimes_{\mathbb{Z}} \mathbb{R}$. Since the normalizer $\mathbf{N}_{\mathbb{C}} := N_{\mathrm{GL}_N(\mathbb{C})}(\mathbf{T}_{\mathbb{C}})$ fixes $\mathrm{Char}(f)$, $\mathbf{N}_{\mathbb{C}}$ is isometric on the metric space $\Gamma \otimes_{\mathbb{Z}} \mathbb{R}$ induced by $\langle\ ,\ \rangle$. Since the Weyl group W is a subgroup of $\mathbf{N}_{\mathbb{C}}$, we conclude that W is isometric on $\Gamma \otimes_{\mathbb{Z}} \mathbb{R}$.

Consider the set of almost simple factors of $\mathbf{G}_{\mathbb{C}}$: $\mathbf{H}_{j,\mathbb{C}}$ for all $1 \le j \le m$. Then $\mathbf{T}_{j,\mathbb{C}} := (\mathbf{T}_{\mathbb{C}} \cap \mathbf{H}_{j,\mathbb{C}})^\circ$ is a maximal torus of $\mathbf{H}_{j,\mathbb{C}}$. Let Γ_j be the character group of $\mathbf{T}_{j,\mathbb{C}}$ and R_j (resp. W_j) the roots (resp. Weyl group) of $\mathbf{H}_{j,\mathbb{C}}$ with respect to the maximal torus $\mathbf{T}_{j,\mathbb{C}}$. Since R_j spans $\Gamma_j \otimes_{\mathbb{Z}} \mathbb{R}$, the disjoint union $R = \coprod_{j=1}^m R_j$ induces a direct sum decomposition:

$$\Gamma \otimes_{\mathbb{Z}} \mathbb{R} = \bigoplus_{j=1}^{m} \Gamma_j \otimes_{\mathbb{Z}} \mathbb{R}. \tag{27}$$

As $W \cong \prod_{j=1}^m W_j$ is isometric on $\Gamma \otimes_{\mathbb{Z}} \mathbb{R}$ and preserves $\Gamma_j \otimes_{\mathbb{Z}} \mathbb{R}$ for all $1 \le j \le m$, (27) is an orthogonal direct sum. This implies by [Bo81, VI §1 Prop. 5 Cor. (i)] that for $1 \le j \le m$,

$$\langle\ ,\ \rangle|_{\Gamma_j \otimes_{\mathbb{Z}} \mathbb{R}} \text{ is a positive multiple of the } \textit{Killing form} \text{ [FH91, §14] on } \Gamma_j \otimes_{\mathbb{Z}} \mathbb{R}. \tag{28}$$

Suppose $\mathbf{G}'_{\mathbb{C}}$ is another connected complex semisimple Lie subgroup of $\mathrm{GL}_N(\mathbb{C})$ sharing the same maximal torus $\mathbf{T}_{\mathbb{C}}$ with $\mathbf{G}_{\mathbb{C}}$. Let R' be the roots of $\mathbf{G}'_{\mathbb{C}}$ with respect to $\mathbf{T}_{\mathbb{C}}$. Then $R' \subset \Gamma \subset \Gamma \otimes_{\mathbb{Z}} \mathbb{R}$. There is a strong geometric connection between R and R'. Let $r \in R$, $r' \in R'$, and θ the angle between them. Since r' and r are characters of $\mathbf{T}_{\mathbb{C}}$, we obtain

$$2\frac{\langle r', r\rangle}{\langle r, r\rangle} \in \mathbb{Z}, \quad 2\frac{\langle r, r'\rangle}{\langle r', r'\rangle} \in \mathbb{Z} \tag{29}$$

by (28). This implies $4\cos^2\theta \in \mathbb{Z}$ and θ can only be a multiple of 30° or 45° (see [Hui13, §2]).

Example: Notation as above. The semisimple Lie group $\mathbf{G}_{\mathbb{C}} = \mathrm{SL}_2(\mathbb{C}) \times \mathrm{SL}_2(\mathbb{C})/\pm(-\mathrm{Id}, -\mathrm{Id})$ is a subgroup of the simple exceptional Lie group $\mathbf{G}'_{\mathbb{C}}$ of type G_2 and f is the restriction to $\mathbf{G}_{\mathbb{C}}$ of the adjoint representation $\mathrm{Ad} : \mathbf{G}'_{\mathbb{C}} \hookrightarrow \mathrm{GL}_N(\mathbb{C})$. Since $\mathbf{T}_{\mathbb{C}}$ is also a maximal torus of $\mathbf{G}'_{\mathbb{C}}$, the configuration of the nonzero weights of f on $\Gamma \otimes_{\mathbb{Z}} \mathbb{R}$ with respect to $\langle\ ,\ \rangle$ is just some positive multiple of the root system of G_2 (the figure). As f contains the adjoint representation of $\mathbf{G}_{\mathbb{C}}$ as a subrepresentation, the roots $R = R_1 \coprod R_2$ of $\mathbf{G}_{\mathbb{C}}$ must appear in the figure. Since (27) is an orthogonal sum, R_1 and R_2 are perpendicular. Without loss of generality, assume $R_1 = \{e_1, -e_1\}$ and $R_2 = \{e_2, -e_2\}$. Then we obtain $||e_2|| = \sqrt{3}||e_1||$. On the other hand, we see that the angles between R (red) and R' (red or blue) are multiples of 30°.



3.2. **Proof of Theorem 3.2.** The notation in §3.1 remains in force. There is a lattice Λ of $\Gamma \otimes_{\mathbb{Z}} \mathbb{R}$ containing Γ, known as the *weight lattice*. The construction is as follows. Let $\widetilde{\mathbf{G}_{\mathbb{C}}}$ be the universal covering of $\mathbf{G}_{\mathbb{C}}$, $\widetilde{\mathbf{T}_{\mathbb{C}}}$ a maximal torus of $\widetilde{\mathbf{G}_{\mathbb{C}}}$ mapping onto $\mathbf{T}_{\mathbb{C}}$, and Λ the character group of $\widetilde{\mathbf{T}_{\mathbb{C}}}$. Then the character group Γ of $\mathbf{T}_{\mathbb{C}}$ injects into Λ as a finite index subgroup. Hence, $\Lambda \otimes_{\mathbb{Z}} \mathbb{R}$ can be identified as $\Gamma \otimes_{\mathbb{Z}} \mathbb{R}$. Suppose $\mathbf{G}_{\mathbb{C}}$ is simple and let $g \in \mathbf{N}_{\mathbb{C}}$. Then g is orthogonal on $\Lambda \otimes_{\mathbb{Z}} \mathbb{R}$. Denote by $\mathbf{G}'_{\mathbb{C}}$ the subgroup $g\mathbf{G}_{\mathbb{C}}g^{-1}$ in $\mathrm{GL}_N(\mathbb{C})$. Then $\mathbf{G}_{\mathbb{C}}$ and $\mathbf{G}'_{\mathbb{C}}$ share the common maximal torus $\mathbf{T}_{\mathbb{C}}$ and $R' = gR$. We first prove that $R = R'$ in Λ if the Lie algebra $\mathfrak{g}$ of $\mathbf{G}_{\mathbb{C}}$ is not equal to A_7, A_8, B_4, D_8 (Theorem 3.2(i)). The configuration $R \subset \Lambda \subset \Lambda \otimes_{\mathbb{Z}} \mathbb{R}$, up to normalization, is the same as the configuration of the root system of the complex simple Lie algebra $\mathfrak{g}$ [GOV94, Table 1].

3.2.1. $\mathfrak{g} = A_n$. Let $r \in R$, $r' \in R'$, and θ the angle between them. Since $R' = gR$, we obtain $||r|| = ||r'||$. Thus, θ is a multiple of 60° or 90° ($2\cos\theta \in \mathbb{Z}$ by (29)). We follow the crucial computation in [Hui13, Prop. 2.8]. Normalize $\Lambda \otimes_{\mathbb{Z}} \mathbb{R}$ such that $||r|| = \sqrt{2}$. There exist weights $e_1, ..., e_n, e_{n+1} \in \Lambda$ such that

$$\begin{aligned} 0 &= e_1 + e_2 + ... + e_n + e_{n+1} \\ \Lambda &= \mathbb{Z}e_1 \oplus \cdots \oplus \mathbb{Z}e_n \\ \langle e_i, e_j \rangle &= \begin{cases} \frac{n}{n+1} & \text{if} \quad 1 \le i = j \le n \\ \frac{-1}{n+1} & \text{if} \quad 1 \le i \ne j \le n \end{cases} \end{aligned} \tag{30}$$

and the roots R comprise the set $\{e_i - e_j : 1 \le i \ne j \le n+1\}$. Since $r' \in \Lambda$, write $r' = a_1e_1 + \cdots + a_ne_n$ as an integral combination. Consider $r = e_i - e_j$ where $i, j \le n$. By (30), we have

$$\begin{aligned} 2\cos\theta = \langle r', r \rangle &= \langle a_1e_1 + \cdots + a_ne_n, e_i - e_j \rangle \\ &= \frac{a_i(n+1) - \sum_{k=1}^n a_k}{n+1} - \frac{a_j(n+1) - \sum_{k=1}^n a_k}{n+1} = a_i - a_j. \end{aligned} \tag{31}$$

Then consider $r = e_i - e_{n+1}$ where $i \leq n$. By (30), we have $r = e_1 + e_2 + \cdots + e_{i-1} + 2e_i + e_{i+1} + \cdots + e_n$, so

$$\begin{aligned} 2\cos\theta = \langle r', r\rangle &= \langle a_1e_1 + \cdots + a_ne_n, e_1 + e_2 + \cdots + e_{i-1} + 2e_i + e_{i+1} + \cdots + e_n\rangle \\ &= \frac{a_i(n+1) - \sum_{k=1}^n a_k}{n+1} + \sum_{j=1}^n \frac{a_j(n+1) - \sum_{k=1}^n a_k}{n+1} = a_i. \end{aligned} \tag{32}$$

Suppose $r' \notin R$, then $\theta \in \{60°, 90°, 120°, 240°, 270°, 300°\}$ and

$$|a_i|, |a_i - a_j| \in \{0, 1\}, \ \ \forall 1 \leq i \neq j \leq n \tag{33}$$

by (31) and (32), which means that either all $a_i \in \{0, 1\}$ or all $a_i \in \{0, -1\}$. Therefore, we obtain

$$\begin{aligned} 2 = \langle r', r'\rangle = \langle a_1e_1 + \cdots + a_ne_n, a_1e_1 + \cdots + a_ne_n\rangle &= \frac{\sum_{i=1}^n na_i^2 - 2\sum_{i<j} a_ia_j}{n+1} \\ = \frac{\sum_{i=1}^n a_i^2 + \sum_{i<j}(a_i - a_j)^2}{n+1} &= \frac{(n-k) + k(n-k)}{n+1}, \end{aligned} \tag{34}$$

where k is the number of zero a_i. Since the integral equation (34) has a solution $(k, n) \in \mathbb{Z}_{\geq 0} \times \mathbb{N}$ such that $k \leq n$ (i.e., $r' \notin R$) if and only if $n = 7, 8$ [Hui13, (**2.13**) $\theta = 60°$], we conclude that $R = R'$ if $n \in \mathbb{N}\backslash\{7, 8\}$.

3.2.2. $\mathfrak{g} = B_n$. Under suitable normalization, $\Lambda \otimes_{\mathbb{Z}} \mathbb{R}$ is isometric to the Euclidean space $\mathbb{R}^n$. The weight lattice Λ is the lattice generated by the standard basis $\{e_1, ..., e_n\}$ and $\frac{e_1+\cdots+e_n}{2}$. The roots $R = \{\pm e_i, \pm e_i \pm e_j : \ 1 \leq i \neq j \leq n\}$.

Let $r' \in R'$ be a short root, i.e., $||r'|| = 1$. Write $r' = \frac{a_1e_1+\cdots+a_ne_n}{2}$ as a half-integral combination. Then either all a_i are even or all a_i are odd. In the even case, $||r'|| = 1$ implies $r' \in R$. In the odd case, $||r'|| = 1$ implies $n = 4$. Hence, the short roots R'° of R' belong to R whenever $n \neq 4$. Since R'° determines R', we obtain $R = R'$ if $n \neq 4$.

3.2.3. $\mathfrak{g} = C_n$. Under suitable normalization, $\Lambda \otimes_{\mathbb{Z}} \mathbb{R}$ is isometric to the Euclidean space $\mathbb{R}^n$. The weight lattice Λ is the lattice generated by the standard basis $e_1, ..., e_n$. The roots $R = \{\pm 2e_i, \pm e_i \pm e_j : \ 1 \leq i \neq j \leq n\}$.

Since all weights of length $\sqrt{2}$ belong to R, the short roots R'° of R' belong to R. Since R'° determines R', we obtain $R = R'$.

3.2.4. $\mathfrak{g} = D_n$. Under suitable normalization, $\Lambda \otimes_{\mathbb{Z}} \mathbb{R}$ is isometric to the Euclidean space $\mathbb{R}^n$. The weight lattice Λ is the lattice generated by the standard basis $e_1, ..., e_n$ and $\frac{e_1+\cdots+e_n}{2}$ (same as B_n). The roots $R = \{\pm e_i \pm e_j : \ 1 \leq i \neq j \leq n\}$.

Let $r' \in R'$ be a root, $||r'|| = \sqrt{2}$. Write $r' = \frac{a_1e_1+\cdots+a_ne_n}{2}$ as a half-integral combination. Then either all a_i are even or all a_i are odd. In the even case, $||r'|| = \sqrt{2}$ implies $r' \in R$. In the odd case, $||r'|| = \sqrt{2}$ implies $n = 8$. This implies $R = R'$ if $n \neq 8$.

3.2.5. $\mathfrak{g} = E_6, E_7, E_8$. Denote the weight lattice (resp. the roots) of E_6, E_7, E_8 by $\Lambda_6, \Lambda_7, \Lambda_8$ (resp. R_6,R_7,R_8).

E_8: Under suitable normalization, $\Lambda_8 \otimes_{\mathbb{Z}} \mathbb{R}$ is isometric to the Euclidean space $\mathbb{R}^8$. Let $e_1, e_2, ..., e_8$ be the standard basis of $\mathbb{R}^8$. The weight lattice Λ_8 is the well-known E_8-lattice (even coordinate system) [CS88, §8.1]

$$\Lambda_8 := \{w = \sum_{i=1}^{8} a_i e_i \in \mathbb{Z}^8 \cup (\mathbb{Z} + 1/2)^8 : \ \sum_{i=1}^{8} a_i \equiv 0 \bmod 2\}, \tag{35}$$

where R_8 consists of all the 240 weights of length $\sqrt{2}$, e.g., $e_1 - e_2$ and $e_2 - e_3$. This implies $R = R'$.

E_7: Under suitable normalization, $\Lambda_7 \otimes_{\mathbb{Z}} \mathbb{R}$ is the orthogonal complement in $\Lambda_8 \otimes_{\mathbb{Z}} \mathbb{R}$ of any root $\alpha \in R_8$. The set of roots R_7 is just the set of 126 elements in $R_8 \cap (\Lambda_7 \otimes_{\mathbb{Z}} \mathbb{R})$ and Λ_7 is the orthogonal projection of Λ_8 to $\Lambda_7 \otimes_{\mathbb{Z}} \mathbb{R}$. To see the last point, let $\mathbf{G}'_{\mathbb{C}}$ be a complex simple Lie group of type E_8, it contains a subgroup $\mathrm{SL}_2(\mathbb{C})$. By [Ad96, Ch. 8], the identity component of the centralizer of $\mathrm{SL}_2(\mathbb{C})$ in $\mathbf{G}'_{\mathbb{C}}$, denoted by $\mathbf{G}_{\mathbb{C}}$, is of type E_7 and contains the center of $\mathrm{SL}_2(\mathbb{C})$. This implies $\mathbf{G}_{\mathbb{C}}$ is simply connected. As $\mathbf{G}'_{\mathbb{C}}$ is also simply connected, the embedding $\mathbf{G}_{\mathbb{C}} \hookrightarrow \mathbf{G}'_{\mathbb{C}}$ induces a surjection map of the weight lattices $\Lambda_8 \twoheadrightarrow \Lambda_7$. This is an explicit description of Λ_7. For example, we take $\alpha = e_1 - e_2$. If $w = a_1 e_1 + \cdots + a_8 e_8 \in \Lambda_8$, then the orthogonal projection along α is

$$w' := (\frac{a_1 + a_2}{2})e_1 + (\frac{a_1 + a_2}{2})e_2 + a_3 e_3 + \cdots + a_8 e_8. \tag{36}$$

Suppose $||w'|| = \sqrt{2}$ and the a_i are integers, then $a_1 + a_2$ must be even for the integral equation $||w'||^2 = 2$ and we see by (36),(35) that $w' \in \Lambda_8$. Thus, $w' \in R_8 \cap (\Lambda_7 \otimes_{\mathbb{Z}} \mathbb{R}) = R_7$. Suppose $||w'|| = \sqrt{2}$ and all the a_i are half integers, write $a_i = b_i/2$. We have two cases. First, if $b_1 + b_2$ is even, then $w' \in \Lambda_8$ by (36),(35). Thus, $w' \in R_8 \cap (\Lambda_7 \otimes_{\mathbb{Z}} \mathbb{R}) = R_7$. Second, if $b_1 + b_2$ is odd, then $||w'||^2 = 2$ is equivalent to

$$(b_1 + b_2)^2 + 2(b_3^2 + \cdots + b_8^2) = 16, \tag{37}$$

which is impossible. Therefore, any weight $w' \in \Lambda_7$ of length $\sqrt{2}$ belongs to R_7. We conclude that $R = R'$.

E_6: The strategy is similar to E_7. Under suitable normalization, $\Lambda_6 \otimes_{\mathbb{Z}} \mathbb{R}$ is the orthogonal complement in $\Lambda_8 \otimes_{\mathbb{Z}} \mathbb{R}$ of a suitable chosen pair of roots $\alpha, \beta \in R_8$. The set of roots R_6 is just the set of 72 elements in $R_8 \cap (\Lambda_6 \otimes_{\mathbb{Z}} \mathbb{R})$ and Λ_6 is the orthogonal projection of Λ_8 to $\Lambda_6 \otimes_{\mathbb{Z}} \mathbb{R}$. To see the last point, let $\mathbf{G}'_{\mathbb{C}}$ be a complex simple Lie group of type E_8, it contains a subgroup $\mathrm{SL}_3(\mathbb{C})$. By [Ad96, Ch. 8], the identity component of the centralizer of $\mathrm{SL}_3(\mathbb{C})$ in $\mathbf{G}'_{\mathbb{C}}$, denoted by $\mathbf{G}_{\mathbb{C}}$, is of type E_6 and contains the center of $\mathrm{SL}_3(\mathbb{C})$. This implies $\mathbf{G}_{\mathbb{C}}$ is simply connected. As $\mathbf{G}'_{\mathbb{C}}$ is also simply connected, the embedding $\mathbf{G}_{\mathbb{C}} \hookrightarrow \mathbf{G}'_{\mathbb{C}}$ induces a surjection map of the weight lattices $\Lambda_8 \twoheadrightarrow \Lambda_6$. This is an explicit description of Λ_6. For example, we take $\alpha = e_1 - e_2, \beta = e_2 - e_3$. If

$w = a_1e_1 + \cdots + a_8e_8 \in \Lambda_8$, then the orthogonal projection along α, β is

$$w' := (\frac{a_1+a_2+a_3}{3})e_1 + (\frac{a_1+a_2+a_3}{3})e_2 + (\frac{a_1+a_2+a_3}{3})e_3 + a_4e_4 + \cdots + a_8e_8. \tag{38}$$

Suppose $||w'|| = \sqrt{2}$ and the a_i are integers, then $a_1+a_2+a_3$ must be divisible by 3 for the integral equation $||w'||^2 = 2$ and we see by (38),(35) that $w' \in \Lambda_8$. Thus, $w' \in R_8 \cap (\Lambda_6 \otimes_{\mathbb{Z}} \mathbb{R}) = R_6$. Suppose $||w'|| = \sqrt{2}$ and all the a_i are half integers, write $a_i = b_i/2$. We have two cases. First, if $b_1+b_2+b_3$ is divisible by 3, then $w' \in \Lambda_8$ by (38),(35). Thus, $w' \in R_8 \cap (\Lambda_6 \otimes_{\mathbb{Z}} \mathbb{R}) = R_6$. Second, if $b_1+b_2+b_3$ is not divisible by 3, then $||w'||^2 = 2$ is equivalent to

$$(b_1+b_2+b_3)^2 + 3(b_4^2 + \cdots + b_8^2) = 24, \tag{39}$$

which is impossible. Therefore, any weight $w' \in \Lambda_6$ of length $\sqrt{2}$ belongs to R_6. We conclude that $R = R'$.

3.2.6. $\mathfrak{g} = F_4$. Under suitable normalization, $\Lambda \otimes_{\mathbb{Z}} \mathbb{R}$ is isometric to the Euclidean space $\mathbb{R}^4$. The weight lattice Λ is the lattice generated by the standard basis e_1, e_2, e_3, e_4 and $\frac{e_1+e_2+e_3+e_4}{2}$ (same as B_4). The roots $R = \{\pm e_i, \pm e_i \pm e_j, \frac{\pm e_1 \pm e_2 \pm e_3 \pm e_4}{2} : \ 1 \le i \ne j \le n\}$.

Since R consists of all weights of length 1 or $\sqrt{2}$, we obtain $R = R'$.

3.2.7. $\mathfrak{g} = G_2$. The weight lattice Λ is generated by the roots R. By observing the root system of G_2, one sees that the shortest nonzero weights of Λ belong to R. Hence, the short roots R'° of R' belong to R. Since R'° determines R', we obtain $R = R'$.

3.2.8. *Theorem 3.2(ii).* It remains to prove Theorem 3.2(ii). Let $\mathbf{G}'_{\mathbb{C}}$ be a connected complex simple Lie group with Lie algebra $\mathfrak{g}' \in \{E_7, E_8, F_4\}$. Embed $\mathbf{G}'_{\mathbb{C}}$ in some $\mathrm{GL}_N(\mathbb{C})$. When $\mathbf{G}'_{\mathbb{C}}$ is of type E_7, it contains an equal rank subgroup $\mathbf{G}_{\mathbb{C}}$ of type A_7; when $\mathbf{G}'_{\mathbb{C}}$ is of type E_8, it contains an equal rank subgroup $\mathbf{G}_{\mathbb{C}}$ of type A_8 (resp. D_8); when $\mathbf{G}'_{\mathbb{C}}$ is of type F_4, it contains an equal rank subgroup $\mathbf{G}_{\mathbb{C}}$ of type B_4 [GOV94, Table 5]. Then we obtain an embedding $f : \mathbf{G}_{\mathbb{C}} \hookrightarrow \mathbf{G}'_{\mathbb{C}} \hookrightarrow \mathrm{GL}_N(\mathbb{C})$.

Pick a maximal torus $\mathbf{T}_{\mathbb{C}}$ of $\mathbf{G}_{\mathbb{C}}$, then it is also a maximal torus of $\mathbf{G}'_{\mathbb{C}}$. If the normalizer $\mathbf{N}_{\mathbb{C}} := N_{\mathrm{GL}_N(\mathbb{C})}(\mathbf{T}_{\mathbb{C}})$ preserves the roots R of $\mathbf{G}_{\mathbb{C}}$, then the Weyl group W' of $\mathbf{G}'_{\mathbb{C}}$ preserves R. Since W' is orthogonal on $\Lambda \otimes_{\mathbb{Z}} \mathbb{R}$, W' acts on the set P of *Weyl chambers* of $R \subset \Lambda \otimes_{\mathbb{Z}} \mathbb{R}$ [FH91, §14]. Each Weyl chamber corresponds to a set of positive roots R^+ of R. The order of P is equal to the order of W, the Weyl group of $\mathbf{G}_{\mathbb{C}}$.

Let $R^+ \in P$ and H the stabilizer of R^+ in W'. Since W' is faithful and orthogonal on $\Lambda \otimes_{\mathbb{Z}} \mathbb{R}$, H is an automorphism group of the *Dynkin diagram* [GOV94, Table 1] of $\mathbf{G}_{\mathbb{C}}$ (or $\mathfrak{g}$). As W is simply transitive on P, we obtain $|H| = |W'|/|W|$.

$\mathfrak{g}'$	$\lvert W' \rvert$
E_7	2903040
E_8	696729600
F_4	1152

$\mathfrak{g}$	$\lvert W \rvert$	Dynkin diagram
A_7	$8! = 40320$	
A_8	$9! = 362880$	
B_4	$4! \times 2^4 = 384$	
D_8	$8! \times 2^7 = 5160960$	

For any embedding $f : \mathbf{G}_{\mathbb{C}} \hookrightarrow \mathrm{GL}_N(\mathbb{C})$ in the first paragraph, we have $|H| > 2$ by the table above. This contradicts that the automorphism group of the Dynkin diagram of A_7, A_8, B_4, D_8 has order at most 2. We conclude that $\mathbf{N}_{\mathbb{C}}$ does not preserve the roots R.

Remark 3.3. *The exceptional complex simple Lie group $\mathbf{G}'_{\mathbb{C}}$ of type G_2 contains $\mathrm{SL}_3(\mathbb{C})$ (type A_2) as an equal rank subgroup. The arguments above do not contradict Theorem 3.2(i) for $\mathbf{G}_{\mathbb{C}} = \mathrm{SL}_3(\mathbb{C})$ because the orders of the Weyl groups of G_2 and A_2 are respectively 12 and 6 and $\frac{12}{6} = 2$ is the order of the automorphism of the Dynkin diagram of A_2.*

Acknowledgments

I would like to thank Gabor Wiese for his interests and comments on the paper.

References

[Ad96] Adams, J.F.: *Lectures on exceptional Lie groups*, Chicago Lectures in Mathematics, University of Chicago Press, 1996. 18

[BLGGT14] Barnet-Lamb, T.; Gee, T.; Geraghty, D.; Taylor, R.: Potential automorphy and change of weight, *Ann. of Math.*, **179** (2014), Issue 2, p. 501-609. 2

[Bo81] Bourbaki, N.: *Groupes et algébres de Lie*, Paris: Masson 1981. 15

[Chi92] Chi, Wen Chen: ℓ-adic and λ-adic representations associated to abelian varieties defined over number fields, *Amer. J. Math.* **114** (1992) 315–353. 4, 6

[Chin04] Chin, Chee Whye: Independence of ℓ of monodromy groups, *J.A.M.S.*, Volume **17**, Number 3, 723-747. 4

[Cl90] Clozel, Laurent: *Motifs et formes automorphes: applications du principe de fonctorialité*, in "Automorphic forms, Shimura varieties and L-functions I", Academic Press (1990). 2

[CS88] Conway, J.H.; Sloane, N.J.A.: *Sphere packings, lattices and groups*, New York: Springer-Verlag, 1988. 18

[De74] Deligne, Pierre: La conjecture de Weil I, *Publ. Math. I.H.E.S.*, **43** (1974), 273-307. 2

[De80] Deligne, Pierre: La conjecture de Weil II, *Publ. Math. I.H.E.S.*, **52** (1980), 138-252. 6

[DMOS82] Deligne, P.; Milne, J.S.; Ogus, A.; Shih, K.Y.: *Hodge cycles, motives, and Shimura varieties*, volume **900** of Lecture Notes in Mathematics, Springer-Verlag, Berlin, 1982. 3

[Fa83] Faltings, Gerd: Endlichkeitssätze für abelsche Varietäten über Zahlkörpern. *Invent. Math.* **73** (1983), no. 3, 349–366. 4

[Fa88] Faltings, Gerd: p-adic Hodge theory, *J.A.M.S.*, **1** (1): 255-299. 12

[FM93] Fontaine, Jean-Marc; Mazur, Barry: *Geometric Galois representations*, in Elliptic curves, modular forms and Fermat's last theorem (J.Coates and S.-T.Yau eds.), International Press (1995). 3

[FH91] Fulton, William; Harris, Joe: *Representation Theory*, Graduate Texts in Mathematics 129 (1st ed.), Springer-Verlag 1991. 15, 19

[GOV94] Gorbatsevich, V.V.; Onishchik, A.L.; Vinberg, E.B.: *Lie Groups and Lie Algebras III*, English transl. in Encycl. Math Sc. 41, Springer-Verlag, Berlin, Heidelberg, 1994. 16, 19

[He80] Henniart, Guy: *Représentations ℓ-adiques abéliennes*, in: Bertin, M.-J. (ed.) Séminaires de Théorie des Nombres, Paris 1980-81, pp. 107-126. Boston Basel Stuttgart: Birkhäuser 1982. 8

[Hui13] Hui, Chun Yin: Monodromy of Galois representations and equal-rank subalgebra equivalence, *Math. Res. Lett.* **20** (2013), no. 4, 705-728. 4, 5, 10, 15, 16, 17

[Hui15a] Hui, Chun Yin: ℓ-independence for compatible systems of (mod ℓ) Galois representations, *Compos. Math.*, Vol. **151**, No. 7 (2015), 1215-1241. 4

[Hui15b] Hui, Chun Yin: On rationality of certain type A Galois representations, arXiv:1411.6280. 4, 5, 9, 11, 13, 15

[Kl72] Kleiman, Steven: *Motives*, Algebraic Geometry, Oslo, 1970 (F. Oort, ed.), Walters-Noordhoff, Groningen, 1972, p. 53-82. 3

[Laf02] Lafforgue, Laurent: Chtoucas de Drinfeld et correspondance de Langlands, *Invent. Math.*, **147** (1), p. 1-241 (2002). 2

[Lan79] Langlands, R. P.: *Automorphic representations, Shimura varieties and motives. Ein Märchen*, Automorphic forms, representations and L-functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, RI, 1979, pp. 205–246. 2

[LP90] Larsen, Michael; Pink, Richard: Determining representations from invariant dimensions, *Invent. Math.* **102**, 377-398 (1990). 15

[LP92] Larsen, Michael; Pink, Richard: On ℓ-independence of algebraic monodromy groups in compatible systems of representations, *Invent. Math.* **107** (1992), 603-636. 4

[LP95] Larsen, Michael; Pink, Richard: Abelian varieties, l-adic representations, and l-independence, *Math. Ann.* **302** (1995), no. 3, 561-579. 4

[LP97] Larsen, Michael; Pink, Richard: A connectedness criterion for ℓ-adic Galois representations, *Israel J. Math.* **97** (1997), 1-10. 6

[Pi98] Pink, Richard: ℓ-adic algebraic monodromy groups, cocharacters, and the Mumford-Tate conjecture, *J. Reine Angew. Math.* **495** (1998), 187-237. 4

[Ri76] Ribet, Kenneth A.: Galois action on division points of Abelian varieties with real multiplications, *Amer. J. Math.* **98** (1976), no. 3, 751–804. 4, 7

[Sc94] Schappacher, Norbert: *CM motives and the Taniyama group*, Motives, Proc. Sympos. Pure Math., vol. **55**, Amer. Math. Soc., Providence, RI, 1994, pp. 485–508. 3

[Se67] Serre, Jean-Pierre: *Sur les groupes de Galois attachés aux groupes p-divisibles*, Proceed. Conf. on Local Fields, Springer-Verlag, 1967, p. 113-131. 8

[Se72] Serre, Jean-Pierre: Propriétés galoisiennes des points dórdre fini des courbes elliptiques, *Invent. Math.* **15** (1972), no. 4, 259–331. 4

[Se81] Serre, Jean-Pierre: Letter to K. A. Ribet, Jan. 1, 1981, reproduced in *Coll. Papers*, vol. IV, no. 133. 6

[Se85] Serre, Jean-Pierre: Letter to J. Tate, Jan. 2, 1985. 4

[Se94] Serre, Jean-Pierre: *Propriétés conjecturales des groupes de Galois motiviques et des représentations l-adiques*, Motives (Seattle, WA, 1991), 377–400, Proc. Sympos. Pure Math., 55, Part 1, Amer. Math. Soc., Providence, RI, 1994. 4

[Se98] Serre, Jean-Pierre: *Abelian l-adic representation and elliptic curves*, Research Notes in Mathematics Vol. **7** (2nd ed.), *A K Peters* (1998). 3, 6, 7, 8, 9, 10, 13

[Ta65] Tate, John: *Algebraic cycles and poles of zeta functions*, in Arithmetical Algebraic Geometry, Harper and Row, New York (1965) 93-110. 3

[Ta67] Tate, John: *p-divisible groups*, Proceed. Conf. on Local Fields, Springer-Verlag, 1967, p. 158-183. 9

[Tay04] Taylor, Richard: Galois representations, *Ann. Fac. Sci. Toulouse Math.* (6) **13** (2004), no. 1, p. 73-119. 2

[Wa81] Waldschmidt, Michel: Transcendance et exponentielles en plusieurs variables, *Invent. Math.* **63**, 97-127 (1981). 8